

## THE RIGHT RIEAMAN-LIOUVILLE FRACTIONAL HERMITE-HADAMARD TYPE INEQUALITIES FOR CONVEX FUNCTIONS

MEHMET KUNT, DÜNYA KARAPINAR, SERCAN TURHAN, İMDAT İŞCAN

ABSTRACT. In this paper, with a new approach, a new fractional Hermite-Hadamard type inequality for convex functions is obtained by using only the right Riemann-Liouville fractional integral. Also, to have new fractional trapezoid and midpoint type inequalities for the differentiable convex functions, two new equalities are proved. Our results generalize the studies [1, 4, 6]. We expect that this study will be lead to the new fractional integration studies for Hermite-Hadamard type inequalities.

### 1. INTRODUCTION

Let  $f : I \subseteq \mathbb{R} \rightarrow \mathbb{R}$  be a convex function defined on the interval  $I$  of real numbers and  $a, b \in I$  with  $a < b$ . The inequality

$$f\left(\frac{a+b}{2}\right) \leq \frac{1}{b-a} \int_a^b f(x) dx \leq \frac{f(a)+f(b)}{2} \quad (1.1)$$

is well known in the literature as Hermite-Hadamard's inequalities [2, 3].

In [1, 6], the authors used the following equality to obtain trapezoid type inequalities and some applications:

**Lemma 1.1.** *Let  $a, b \in I$  with  $a < b$  and  $f : I^\circ \rightarrow \mathbb{R}$  is a differentiable mapping ( $I^\circ$  is the interior of  $I$ ). If  $f' \in L[a, b]$ , then we have*

$$\frac{f(a)+f(b)}{2} - \frac{1}{b-a} \int_a^b f(u) du = \frac{b-a}{2} \int_0^1 (1-2t) f'(ta+(1-t)b) dt. \quad (1.2)$$

In [4], Kırmacı used the following equality to obtain midpoint type inequalities and some applications:

---

2000 *Mathematics Subject Classification.* 26A51, 26A33, 26D10.  
*Key words and phrases.* Convex functions, Hermite-Hadamard inequality, Right Rieaman-Liouville fractional integral, Trapezoid type inequalities, Midpoint type inequalities.  
©2018 Ilirias Research Institute, Prishtinë, Kosovë.  
Submitted October 8, 2017. Published December 28, 2017.  
Communicated by Serkan Araci.

**Lemma 1.2.** Let  $a, b \in I$  with  $a < b$  and  $f : I^\circ \rightarrow \mathbb{R}$  is a differentiable mapping ( $I^\circ$  the interior of  $I$ ). If  $f' \in L[a, b]$ , then we have

$$\begin{aligned} & \frac{1}{b-a} \int_a^b f(u) du - f\left(\frac{a+b}{2}\right) \\ &= (b-a) \left[ \int_0^{1/2} t f'(ta + (1-t)b) dt + \int_{1/2}^1 (t-1) f'(ta + (1-t)b) dt \right]. \end{aligned} \quad (1.3)$$

**Definition 1.** [7, page 12]. A function  $f$  defined on  $I$  has a support at  $x_0 \in I$  if there exists an affine functions  $A(x) = f(x_0) + m(x - x_0)$  such that  $A(x) \leq f(x)$  for all  $x \in I$ . The graph of the support function  $A$  is called a line of support for  $f$  at  $x_0$ .

**Theorem 1.3.** [7, page 12]  $f : (a, b) \rightarrow \mathbb{R}$  is a convex function if and only if there is at least one line of support for  $f$  at each  $x_0 \in (a, b)$ .

Following definitions of the left and right side Riemann-Liouville fractional integrals are well known in the literature.

**Definition 2.** Let  $a, b \in \mathbb{R}$  with  $a < b$  and  $f \in L[a, b]$ . The left and right Riemann-Liouville fractional integrals  $J_{a+}^\alpha f$  and  $J_{b-}^\alpha f$  of order  $\alpha > 0$  are defined by

$$J_{a+}^\alpha f(x) = \frac{1}{\Gamma(\alpha)} \int_a^x (x-t)^{\alpha-1} f(t) dt, \quad x > a$$

and

$$J_{b-}^\alpha f(x) = \frac{1}{\Gamma(\alpha)} \int_x^b (t-x)^{\alpha-1} f(t) dt, \quad x < b$$

respectively, where  $\Gamma(\alpha)$  is the Gamma function defined by  $\Gamma(\alpha) = \int_0^\infty e^{-t} t^{\alpha-1} dt$  (see [5, page 69] and [9, page 4]).

In [8], Sarıkaya et al. proved the following Hermite-Hadamard type fractional integral inequalities:

**Theorem 1.4.** Let  $f : [a, b] \rightarrow \mathbb{R}$  be a positive function with  $0 \leq a < b$  and  $f \in L[a, b]$ . If  $f$  is a convex function on  $[a, b]$ , then the following inequality for fractional integrals hold:

$$f\left(\frac{a+b}{2}\right) \leq \frac{\Gamma(\alpha+1)}{2(b-a)^\alpha} [J_{a+}^\alpha f(b) + J_{b-}^\alpha f(a)] \leq \frac{f(a) + f(b)}{2} \quad (1.4)$$

with  $\alpha > 0$ .

**Remark.** In Theorem 1.4, it is not necessary supposing that  $f$  be a positive function and  $a, b$  are positive real numbers. From the Definition 2, it is clear that  $a, b$  are any real numbers such as  $a < b$ .

In literature, there are so many studies for Hermite-Hadamard type inequalities by using the left and right fractional integrals (such as Riemann-Liouville fractional integrals, Hadamard fractional integrals, Conformable fractional integrals etc.). In all of them, the left and right fractional integrals are used together. As much as we know, there is not any study for Hermite-Hadamard type inequalities by using only the right fractional integrals or the left fractional integrals.

In this paper, our aim is to obtain new fractional Hermite-Hadamard type inequalities by using only the right Riemann-Liouville fractional integral for convex functions. Also, to have new fractional trapezoid and midpoint type inequalities, we prove two new identities for the differentiable convex functions. We expect that, this study will be fundamental to the new fractional integration studies for Hermite-Hadamard type inequalities.

## 2. THE RIGHT FRACTIONAL HERMITE HADAMARD INEQUALITY

**Theorem 2.1.** *Let  $a, b \in \mathbb{R}$  with  $a < b$  and  $f : [a, b] \rightarrow \mathbb{R}$  be a convex function. If  $f \in L[a, b]$ , then the following inequality for the right Riemann-Liouville fractional integral holds:*

$$f\left(\frac{a + \alpha b}{\alpha + 1}\right) \leq \frac{\Gamma(\alpha + 1)}{(b - a)^\alpha} J_{b-}^\alpha f(a) \leq \frac{f(a) + \alpha f(b)}{\alpha + 1} \quad (2.1)$$

with  $\alpha > 0$ .

*Proof.* Let  $\alpha > 0$ . Since  $f$  is convex on  $[a, b]$ , using Theorem 1.3, there is at least one line of support

$$A(x) = f\left(\frac{a + \alpha b}{\alpha + 1}\right) + m\left(x - \frac{a + \alpha b}{\alpha + 1}\right) \leq f(x) \quad (2.2)$$

for all  $x \in [a, b]$ . From (2.2), we have

$$\begin{aligned} A(tb + (1 - t)a) &= f\left(\frac{a + \alpha b}{\alpha + 1}\right) + m\left(tb + (1 - t)a - \frac{a + \alpha b}{\alpha + 1}\right) \\ &\leq f(tb + (1 - t)a) \end{aligned} \quad (2.3)$$

for all  $t \in [0, 1]$ . Multiplying both sides of (2.3) with  $\alpha t^{\alpha-1}$  and integrating over  $[0, 1]$  respect to  $t$ , we have

$$\begin{aligned} &\alpha \int_0^1 t^{\alpha-1} A(tb + (1 - t)a) dt \\ &= \alpha \int_0^1 t^{\alpha-1} \left[ f\left(\frac{a + \alpha b}{\alpha + 1}\right) + m\left(tb + (1 - t)a - \frac{a + \alpha b}{\alpha + 1}\right) \right] dt \\ &= \alpha f\left(\frac{a + \alpha b}{\alpha + 1}\right) \int_0^1 t^{\alpha-1} dt + m \left[ \int_0^1 \alpha [t^\alpha b + (t^{\alpha-1} - t^\alpha) a] dt \right. \\ &\quad \left. - \frac{a + \alpha b}{\alpha + 1} \alpha \int_0^1 t^{\alpha-1} dt \right] \\ &= f\left(\frac{a + \alpha b}{\alpha + 1}\right) + m \left[ \frac{a + \alpha b}{\alpha + 1} - \frac{a + \alpha b}{\alpha + 1} \right] \\ &= f\left(\frac{a + \alpha b}{\alpha + 1}\right) \\ &\leq \alpha \int_0^1 t^{\alpha-1} f(tb + (1 - t)a) dt \\ &= \frac{\alpha}{(b - a)^\alpha} \int_a^b (t - a)^{\alpha-1} f(t) dt \\ &= \frac{\Gamma(\alpha + 1)}{(b - a)^\alpha} J_{b-}^\alpha f(a). \end{aligned} \quad (2.4)$$

On the other hand, using the convexity of  $f$  on  $[a, b]$ , we have

$$f(tb + (1-t)a) \leq tf(b) + (1-t)f(a) \quad (2.5)$$

for all  $t \in [0, 1]$ . Multiplying both sides of (2.5) with  $\alpha t^{\alpha-1}$  and integrating over  $[0, 1]$  respect to  $t$ , we have

$$\begin{aligned} \alpha \int_0^1 t^{\alpha-1} f(tb + (1-t)a) dt &= \frac{\alpha}{(b-a)^\alpha} \int_a^b (t-a)^{\alpha-1} f(t) dt \\ &= \frac{\Gamma(\alpha+1)}{(b-a)^\alpha} J_{b-}^\alpha f(a) \\ &\leq \alpha f(b) \int_0^1 t^\alpha dt + \alpha f(a) \int_0^1 (t^{\alpha-1} - t^\alpha) dt \\ &= \frac{f(a) + \alpha f(b)}{\alpha + 1}. \end{aligned} \quad (2.6)$$

By using (2.4) and (2.6), we have (2.1). This completes the proof.  $\square$

**Remark.** In Theorem 2.1, if one takes  $\alpha = 1$ , one has (1.1) (Hermite-Hadamard inequalities).

### 3. LEMMAS

In this section we will prove the two identities related to Lemma 1.1 and Lemma 1.2.

**Lemma 3.1.** Let  $a, b \in \mathbb{R}$  with  $a < b$  and  $f : [a, b] \rightarrow \mathbb{R}$  be a differentiable function on  $(a, b)$ . If  $f' \in L[a, b]$ , then the following equality for the right Riemann-Liouville fractional integrals holds:

$$\begin{aligned} &\frac{f(a) + \alpha f(b)}{\alpha + 1} - \frac{\Gamma(\alpha + 1)}{(b-a)^\alpha} J_{b-}^\alpha f(a) \\ &= \frac{b-a}{\alpha + 1} \int_0^1 [(\alpha + 1)(1-t)^\alpha - 1] f'(ta + (1-t)b) dt \end{aligned} \quad (3.1)$$

with  $\alpha > 0$ .

*Proof.* If we apply the partial integration to the right hand side of the equation (3.1), we have

$$\begin{aligned}
& \frac{b-a}{\alpha+1} \int_0^1 [(\alpha+1)(1-t)^\alpha - 1] f'(ta + (1-t)b) dt \\
= & (b-a) \left[ \int_0^1 (1-t)^\alpha f'(ta + (1-t)b) dt - \frac{1}{\alpha+1} \int_0^1 f'(ta + (1-t)b) dt \right] \\
= & (b-a) \left[ \int_0^1 t^\alpha f'(tb + (1-t)a) dt - \frac{1}{\alpha+1} \int_0^1 f'(tb + (1-t)a) dt \right] \\
= & (b-a) \left[ \begin{aligned} & \left. t^\alpha \frac{f(tb+(1-t)a)}{b-a} \right|_0^1 - \alpha \int_0^1 t^{\alpha-1} \frac{f(tb+(1-t)a)}{b-a} dt \\ & - \frac{1}{\alpha+1} \left. \frac{f(tb+(1-t)a)}{b-a} \right|_0^1 \end{aligned} \right] \\
= & \left[ f(b) - \alpha \int_0^1 t^{\alpha-1} f(tb + (1-t)a) dt - \frac{f(b) - f(a)}{\alpha+1} \right] \\
= & \frac{f(a) + \alpha f(b)}{\alpha+1} - \frac{\Gamma(\alpha+1)}{(b-a)^\alpha} J_{b-}^\alpha f(a).
\end{aligned}$$

This completes the proof.  $\square$

**Remark.** In Lemma 3.1, if one takes  $\alpha = 1$ , one has the Lemma 1.1.

**Lemma 3.2.** Let  $a, b \in \mathbb{R}$  with  $a < b$  and  $f : [a, b] \rightarrow \mathbb{R}$  be a differentiable function on  $(a, b)$ . If  $f' \in L[a, b]$ , then the following equality for the right Riemann-Liouville fractional integrals holds:

$$\begin{aligned}
& \frac{\Gamma(\alpha+1)}{(b-a)^\alpha} J_{b-}^\alpha f(a) - f\left(\frac{a+\alpha b}{\alpha+1}\right) \\
= & (b-a) \left[ \begin{aligned} & \int_{\frac{\alpha}{\alpha+1}}^{\frac{\alpha}{\alpha+1}} -t^\alpha f'(tb + (1-t)a) dt \\ & + \int_{\frac{\alpha}{\alpha+1}}^1 (1-t^\alpha) f'(tb + (1-t)a) dt \end{aligned} \right] \quad (3.2)
\end{aligned}$$

with  $\alpha > 0$ .

*Proof.* If we apply the partial integration to the right hand side of the equation (3.2), we have

$$\begin{aligned}
& (b-a) \left[ \int_0^{\frac{\alpha}{\alpha+1}} -t^\alpha f'(tb+(1-t)a) dt + \int_{\frac{\alpha}{\alpha+1}}^1 (1-t^\alpha) f'(tb+(1-t)a) dt \right] \\
= & (b-a) \left[ \int_0^1 -t^\alpha f'(tb+(1-t)a) dt + \int_{\frac{\alpha}{\alpha+1}}^1 f'(tb+(1-t)a) dt \right] \\
= & (b-a) \left[ -t^\alpha \frac{f(tb+(1-t)a)}{b-a} \Big|_0^1 + \alpha \int_0^1 t^{\alpha-1} \frac{f(tb+(1-t)a)}{b-a} \right. \\
& \left. + \frac{f(tb+(1-t)a)}{b-a} \Big|_{\frac{\alpha}{\alpha+1}}^1 \right] \\
= & \left[ -f(b) + \alpha \int_0^1 t^{\alpha-1} f(tb+(1-t)a) dt + \left( f(b) - f\left(\frac{a+\alpha b}{\alpha+1}\right) \right) \right] \\
= & \left[ \alpha \int_0^1 t^{\alpha-1} f(tb+(1-t)a) dt - f\left(\frac{a+\alpha b}{\alpha+1}\right) \right] \\
= & \frac{\Gamma(\alpha+1)}{(b-a)^\alpha} J_{b-}^\alpha f(a) - f\left(\frac{a+\alpha b}{\alpha+1}\right).
\end{aligned}$$

This completes the proof.  $\square$

**Remark.** In Lemma 3.2, if one takes  $\alpha = 1$ , one has the Lemma 1.2.

#### 4. THE RIGHT FRACTIONAL TRAPEZOID AND MIDPOINT TYPE INEQUALITIES

In this section we will obtain some new right Riemann-Liouville fractional trapezoid and midpoint type inequalities by using Lemma 3.1 and Lemma 3.2.

**Theorem 4.1.** Let  $a, b \in \mathbb{R}$  with  $a < b$  and  $f : [a, b] \rightarrow \mathbb{R}$  be a differentiable function on  $(a, b)$ . If  $|f'|$  is convex on  $[a, b]$ , then the following right Riemann-Liouville fractional integral inequality holds:

$$\begin{aligned}
& \left| \frac{f(a) + \alpha f(b)}{\alpha + 1} - \frac{\Gamma(\alpha + 1)}{(b-a)^\alpha} J_{b-}^\alpha f(a) \right| \\
\leq & \frac{b-a}{\alpha+1} \left[ \begin{array}{l} R_1(\alpha) |f'(b)| + R_2(\alpha) |f'(a)| \\ + R_3(\alpha) |f'(b)| + R_4(\alpha) |f'(a)| \end{array} \right] \quad (4.1)
\end{aligned}$$

where

$$\begin{aligned}
R_1(\alpha) &= \int_0^{\frac{1}{\sqrt[\alpha]{\alpha+1}}} (1 - (\alpha+1)t^\alpha) t dt = \frac{\alpha}{2(\alpha+2)(\alpha+1)^{\frac{2}{\alpha}}}, \\
R_2(\alpha) &= \int_0^{\frac{1}{\sqrt[\alpha]{\alpha+1}}} (1 - (\alpha+1)t^\alpha)(1-t) dt \\
&= \frac{\alpha \left( 2(\alpha+2)(\alpha+1)^{\frac{1}{\alpha}-1} - 1 \right)}{2(\alpha+2)(\alpha+1)^{\frac{2}{\alpha}}}, \\
R_3(\alpha) &= \int_{\frac{1}{\sqrt[\alpha]{\alpha+1}}}^1 ((\alpha+1)t^\alpha - 1) t dt = \frac{\alpha \left( 1 + (\alpha+1)^{\frac{2}{\alpha}} \right)}{2(\alpha+2)(\alpha+1)^{\frac{2}{\alpha}}}, \\
R_4(\alpha) &= \int_{\frac{1}{\sqrt[\alpha]{\alpha+1}}}^1 ((\alpha+1)t^\alpha - 1)(1-t) dt \\
&= \frac{\alpha \left( 2(\alpha+2)(\alpha+1)^{\frac{1}{\alpha}-1} - 1 - (\alpha+1)^{\frac{2}{\alpha}} \right)}{2(\alpha+2)(\alpha+1)^{\frac{2}{\alpha}}},
\end{aligned}$$

with  $\alpha > 0$ .

*Proof.* Using Lemma 3.1 and the convexity of  $|f'|$ , we have

$$\begin{aligned}
& \left| \frac{f(a) + \alpha f(b)}{\alpha+1} - \frac{\Gamma(\alpha+1)}{(b-a)^\alpha} J_{b-}^\alpha f(a) \right| \\
& \leq \frac{b-a}{\alpha+1} \int_0^1 |(\alpha+1)(1-t)^\alpha - 1| |f'(ta + (1-t)b)| dt \\
& = \frac{b-a}{\alpha+1} \int_0^1 |(\alpha+1)t^\alpha - 1| |f'(tb + (1-t)a)| dt \\
& \leq \frac{b-a}{\alpha+1} \left[ \int_0^{\frac{1}{\sqrt[\alpha]{\alpha+1}}} (1 - (\alpha+1)t^\alpha) [t|f'(b)| + (1-t)|f'(a)|] dt \right. \\
& \quad \left. + \int_{\frac{1}{\sqrt[\alpha]{\alpha+1}}}^1 ((\alpha+1)t^\alpha - 1) [t|f'(b)| + (1-t)|f'(a)|] dt \right] \\
& \leq \frac{b-a}{\alpha+1} \left[ \frac{\alpha}{2(\alpha+2)(\alpha+1)^{\frac{2}{\alpha}}} |f'(b)| + \frac{\alpha(2(\alpha+2)(\alpha+1)^{\frac{1}{\alpha}-1} - 1)}{2(\alpha+2)(\alpha+1)^{\frac{2}{\alpha}}} |f'(a)| \right. \\
& \quad \left. + \frac{\alpha(1+(\alpha+1)^{\frac{2}{\alpha}})}{2(\alpha+2)(\alpha+1)^{\frac{2}{\alpha}}} |f'(b)| + \frac{\alpha(2(\alpha+2)(\alpha+1)^{\frac{1}{\alpha}-1} - 1 - (\alpha+1)^{\frac{2}{\alpha}})}{2(\alpha+2)(\alpha+1)^{\frac{2}{\alpha}}} |f'(a)| \right].
\end{aligned}$$

This completes the proof.  $\square$

**Remark.** In Theorem 4.1, if one takes  $\alpha = 1$ , one has the inequality proved in [1, Theorem 2.2].

**Theorem 4.2.** Let  $a, b \in \mathbb{R}$  with  $a < b$  and  $f : [a, b] \rightarrow \mathbb{R}$  be a differentiable function on  $(a, b)$ . If  $|f'|^q$  is convex on  $[a, b]$  for  $q \geq 1$ , then the following right Riemann-Liouville fractional integral inequality holds:

$$\begin{aligned}
& \left| \frac{f(a) + \alpha f(b)}{\alpha+1} - \frac{\Gamma(\alpha+1)}{(b-a)^\alpha} J_{b-}^\alpha f(a) \right| \\
& \leq \frac{b-a}{\alpha+1} \left( \frac{2\alpha}{(\alpha+1)^{1+\frac{1}{\alpha}}} \right)^{1-\frac{1}{q}} \left( \begin{array}{l} R_1(\alpha) |f'(b)|^q + R_2(\alpha) |f'(a)|^q \\ + R_3(\alpha) |f'(b)|^q + R_4(\alpha) |f'(a)|^q \end{array} \right)^{\frac{1}{q}} \quad (4.2)
\end{aligned}$$

where  $R_1(\alpha)$ - $R_4(\alpha)$  are the same as in Theorem 4.1 and  $\alpha > 0$ .

*Proof.* Using Lemma 3.1, power mean inequality and the convexity of  $|f'|^q$ , we have

$$\begin{aligned}
& \left| \frac{f(a) + \alpha f(b)}{\alpha + 1} - \frac{\Gamma(\alpha + 1)}{(b-a)^\alpha} J_{b-}^\alpha f(a) \right| \\
& \leq \frac{b-a}{\alpha+1} \int_0^1 |(\alpha+1)(1-t)^\alpha - 1| |f'(ta + (1-t)b)| dt \\
& = \frac{b-a}{\alpha+1} \int_0^1 |(\alpha+1)t^\alpha - 1| |f'(tb + (1-t)a)| dt \\
& \leq \frac{b-a}{\alpha+1} \left[ \begin{aligned} & \left( \int_0^1 |(\alpha+1)t^\alpha - 1| dt \right)^{1-\frac{1}{q}} \\ & \times \left( \int_0^1 |(\alpha+1)t^\alpha - 1| |f'(tb + (1-t)b)|^q dt \right)^{\frac{1}{q}} \end{aligned} \right] \\
& \leq \frac{b-a}{\alpha+1} \left[ \begin{aligned} & \left( \int_0^{\frac{1}{\sqrt[\alpha]{\alpha+1}}} (1 - (\alpha+1)t^\alpha) dt + \int_{\frac{1}{\sqrt[\alpha]{\alpha+1}}}^1 ((\alpha+1)t^\alpha - 1) dt \right)^{1-\frac{1}{q}} \\ & \times \left( \int_0^{\frac{1}{\sqrt[\alpha]{\alpha+1}}} (1 - (\alpha+1)t^\alpha) [t|f'(b)|^q + (1-t)|f'(a)|^q] dt \right. \\ & \left. + \int_{\frac{1}{\sqrt[\alpha]{\alpha+1}}}^1 ((\alpha+1)t^\alpha - 1) [t|f'(b)|^q + (1-t)|f'(a)|^q] dt \right)^{\frac{1}{q}} \end{aligned} \right] \\
& \leq \frac{b-a}{\alpha+1} \left( \frac{2\alpha}{(\alpha+1)^{1+\frac{1}{\alpha}}} \right)^{1-\frac{1}{q}} \left( \begin{aligned} & R_1(\alpha) |f'(b)|^q + R_2(\alpha) |f'(a)|^q \\ & + R_3(\alpha) |f'(b)|^q + T_4(\alpha) |f'(a)|^q \end{aligned} \right)^{\frac{1}{q}}.
\end{aligned}$$

This completes the proof.  $\square$

**Remark.** In Theorem 4.2, if one takes  $\alpha = 1$ , one has the inequality proved in [6, Theorem 1].

**Theorem 4.3.** Let  $a, b \in \mathbb{R}$  with  $a < b$  and  $f : [a, b] \rightarrow \mathbb{R}$  be a differentiable function on  $(a, b)$ . If  $|f'|^q$  is convex on  $[a, b]$  for  $q > 1$ , then the following right Riemann-Liouville fractional integral inequality holds:

$$\begin{aligned}
& \left| \frac{f(a) + \alpha f(b)}{\alpha + 1} - \frac{\Gamma(\alpha + 1)}{(b-a)^\alpha} J_{b-}^\alpha f(a) \right| \\
& \leq \frac{b-a}{\alpha+1} (R_5(\alpha, p) + R_6(\alpha, p))^{\frac{1}{p}} \left( \frac{|f'(b)|^q + |f'(a)|^q}{2} \right)^{\frac{1}{q}} \quad (4.3)
\end{aligned}$$

where

$$\begin{aligned}
R_5(\alpha, p) &= \int_0^{\frac{1}{\sqrt[\alpha]{\alpha+1}}} (1 - (\alpha+1)t^\alpha)^p dt, \\
R_6(\alpha, p) &= \int_{\frac{1}{\sqrt[\alpha]{\alpha+1}}}^1 ((\alpha+1)t^\alpha - 1)^p dt,
\end{aligned}$$

with  $\frac{1}{p} + \frac{1}{q} = 1$  and  $\alpha > 0$ .



*Proof.* Using Lemma 3.1, Hölder inequality and the convexity of  $|f'|^q$ , we have

$$\begin{aligned}
& \left| \frac{f(a) + \alpha f(b)}{\alpha + 1} - \frac{\Gamma(\alpha + 1)}{(b-a)^\alpha} J_{b-}^\alpha f(a) \right| \\
& \leq \frac{b-a}{\alpha+1} \int_0^1 |(\alpha+1)(1-t)^\alpha - 1| |f'(ta + (1-t)b)| dt \\
& = \frac{b-a}{\alpha+1} \int_0^1 |(\alpha+1)t^\alpha - 1| |f'(tb + (1-t)a)| dt \\
& = \frac{b-a}{\alpha+1} \int_0^1 |1 - (\alpha+1)t^\alpha| |f'(tb + (1-t)a)| dt \\
& \leq \frac{b-a}{\alpha+1} \left[ \left( \int_0^1 |1 - (\alpha+1)t^\alpha|^p dt \right)^{\frac{1}{p}} \left( \int_0^1 |f'(tb + (1-t)a)|^q dt \right)^{\frac{1}{q}} \right] \\
& \leq \frac{b-a}{\alpha+1} \left[ \left( \int_0^{\frac{1}{\sqrt[\alpha]{\alpha+1}}} (1 - (\alpha+1)t^\alpha)^p dt + \int_{\frac{1}{\sqrt[\alpha]{\alpha+1}}}^1 ((\alpha+1)t^\alpha - 1)^p dt \right)^{\frac{1}{p}} \right. \\
& \quad \left. \times \left( \int_0^1 [t|f'(b)|^q + (1-t)|f'(a)|^q] dt \right)^{\frac{1}{q}} \right] \\
& \leq \frac{b-a}{\alpha+1} \left[ (R_5(\alpha, p) + R_6(\alpha, p))^{\frac{1}{p}} \left( \frac{|f'(b)|^q + |f'(a)|^q}{2} \right)^{\frac{1}{q}} \right].
\end{aligned}$$

This completes the proof.  $\square$

**Remark.** In Theorem 4.3, if one takes  $\alpha = 1$ , one has the inequality proved in [1, Theorem 2.3].

**Theorem 4.4.** Let  $a, b \in \mathbb{R}$  with  $a < b$  and  $f : [a, b] \rightarrow \mathbb{R}$  be a differentiable function on  $(a, b)$ . If  $|f'|$  is convex on  $[a, b]$ , then the following right Riemann-Liouville fractional integral inequality holds:

$$\begin{aligned}
& \left| \frac{\Gamma(\alpha + 1)}{(b-a)^\alpha} J_{b-}^\alpha f(a) - f\left(\frac{a + \alpha b}{\alpha + 1}\right) \right| \\
& \leq (b-a) \left[ \begin{array}{l} R_7(\alpha) |f'(b)| + R_8(\alpha) |f'(a)| \\ + R_9(\alpha) |f'(b)| + R_{10}(\alpha) |f'(a)| \end{array} \right] \quad (4.4)
\end{aligned}$$

where

$$\begin{aligned}
R_7(\alpha) &= \int_0^{\frac{\alpha}{\alpha+1}} t^{\alpha+1} dt = \frac{\alpha \left(\frac{\alpha}{\alpha+1}\right)^{\alpha+1}}{(\alpha+1)(\alpha+2)}, \\
R_8(\alpha) &= \int_0^{\frac{\alpha}{\alpha+1}} t^\alpha - t^{\alpha+1} dt = \frac{2 \left(\frac{\alpha}{\alpha+1}\right)^{\alpha+1}}{(\alpha+1)(\alpha+2)}, \\
R_9(\alpha) &= \int_{\frac{\alpha}{\alpha+1}}^1 t - t^{\alpha+1} dt = \frac{\alpha(2\alpha^\alpha + (\alpha+1)^\alpha)}{2(\alpha+1)^{\alpha+2}(\alpha+2)}, \\
R_{10}(\alpha) &= \int_{\frac{\alpha}{\alpha+1}}^1 (1-t^\alpha)(1-t) dt = \frac{4\alpha^{\alpha+1} - \alpha(\alpha+1)^\alpha}{2(\alpha+1)^{\alpha+2}(\alpha+2)},
\end{aligned}$$

with  $\alpha > 0$ .

*Proof.* Using Lemma 3.2 and the convexity of  $|f'|$ , we have

$$\begin{aligned}
& \left| \frac{\Gamma(\alpha+1)}{(b-a)^\alpha} J_{b-}^\alpha f(a) - f\left(\frac{a+\alpha b}{\alpha+1}\right) \right| \\
& \leq (b-a) \left[ \int_0^{\frac{\alpha}{\alpha+1}} t^\alpha |f'(tb+(1-t)a)| dt \right. \\
& \quad \left. + \int_{\frac{\alpha}{\alpha+1}}^1 (1-t)^\alpha |f'(tb+(1-t)a)| dt \right] \\
& \leq (b-a) \left[ \int_0^{\frac{\alpha}{\alpha+1}} t^\alpha [t|f'(b)| + (1-t)|f'(a)|] dt \right. \\
& \quad \left. + \int_{\frac{\alpha}{\alpha+1}}^1 (1-t)^\alpha [t|f'(b)| + (1-t)|f'(a)|] dt \right] \\
& \leq (b-a) \left[ \frac{\alpha\left(\frac{\alpha}{\alpha+1}\right)^{\alpha+1}}{(\alpha+1)(\alpha+2)} |f'(b)| + \frac{2\left(\frac{\alpha}{\alpha+1}\right)^{\alpha+1}}{(\alpha+1)(\alpha+2)} |f'(a)| \right. \\
& \quad \left. + \frac{\alpha(2\alpha^\alpha + (\alpha+1)^\alpha)}{2(\alpha+1)^{\alpha+2}(\alpha+2)} |f'(b)| + \frac{4\alpha^{\alpha+1} - \alpha(\alpha+1)^\alpha}{2(\alpha+1)^{\alpha+2}(\alpha+2)} |f'(a)| \right].
\end{aligned}$$

This completes the proof.  $\square$

**Remark.** In Theorem 4.4, if one takes  $\alpha = 1$ , one has the inequality proved in [4, Theorem 2.2].

**Theorem 4.5.** Let  $a, b \in \mathbb{R}$  with  $a < b$  and  $f : [a, b] \rightarrow \mathbb{R}$  be a differentiable function on  $(a, b)$ . If  $|f'|^q$  is convex on  $[a, b]$  for  $q \geq 1$ , then the following right Riemann-Liouville fractional integral inequality holds:

$$\begin{aligned}
& \left| \frac{\Gamma(\alpha+1)}{(b-a)^\alpha} J_{b-}^\alpha f(a) - f\left(\frac{a+\alpha b}{\alpha+1}\right) \right| \\
& \leq (b-a) \left( \frac{\alpha^{\alpha+1}}{(\alpha+1)^{\alpha+2}} \right)^{1-\frac{1}{q}} \left[ \begin{aligned} & (R_7(\alpha) |f'(b)|^q + R_8(\alpha) |f'(a)|^q)^{\frac{1}{q}} \\ & + (R_9(\alpha) |f'(b)|^q + R_{10}(\alpha) |f'(a)|^q)^{\frac{1}{q}} \end{aligned} \right] \quad (4.5)
\end{aligned}$$

where  $R_7(\alpha)$ - $R_{10}(\alpha)$  are the same as in Theorem 4.4 and  $\alpha > 0$ .

*Proof.* Using Lemma 3.2, power mean inequality and the convexity of  $|f'|^q$ , we have

$$\begin{aligned}
& \left| \frac{\Gamma(\alpha+1)}{(b-a)^\alpha} J_{b-}^\alpha f(a) - f\left(\frac{a+\alpha b}{\alpha+1}\right) \right| \\
& \leq (b-a) \left[ \int_0^{\frac{\alpha}{\alpha+1}} t^\alpha |f'(tb+(1-t)a)| dt + \int_{\frac{\alpha}{\alpha+1}}^1 (1-t)^\alpha |f'(tb+(1-t)a)| dt \right] \\
& \leq (b-a) \left[ \left( \int_0^{\frac{\alpha}{\alpha+1}} t^\alpha dt \right)^{1-\frac{1}{q}} \left( \int_0^{\frac{\alpha}{\alpha+1}} t^\alpha |f'(tb+(1-t)a)|^q dt \right)^{\frac{1}{q}} \right. \\
& \quad \left. + \left( \int_{\frac{\alpha}{\alpha+1}}^1 (1-t)^\alpha dt \right)^{1-\frac{1}{q}} \left( \int_{\frac{\alpha}{\alpha+1}}^1 (1-t)^\alpha |f'(tb+(1-t)a)|^q dt \right)^{\frac{1}{q}} \right] \\
& \leq (b-a) \left( \frac{\alpha^{\alpha+1}}{(\alpha+1)^{\alpha+2}} \right)^{1-\frac{1}{q}} \\
& \quad \times \left[ \left( \int_0^{\frac{\alpha}{\alpha+1}} t^\alpha [t|f'(b)|^q + (1-t)|f'(a)|^q] dt \right)^{\frac{1}{q}} \right. \\
& \quad \left. + \left( \int_{\frac{\alpha}{\alpha+1}}^1 (1-t)^\alpha [t|f'(b)|^q + (1-t)|f'(a)|^q] dt \right)^{\frac{1}{q}} \right] \\
& \leq (b-a) \left( \frac{\alpha^{\alpha+1}}{(\alpha+1)^{\alpha+2}} \right)^{1-\frac{1}{q}} \\
& \quad \times \left[ \left( \frac{\alpha \left(\frac{\alpha}{\alpha+1}\right)^{\alpha+1}}{(\alpha+1)(\alpha+2)} |f'(b)|^q + \frac{2 \left(\frac{\alpha}{\alpha+1}\right)^{\alpha+1}}{(\alpha+1)(\alpha+2)} |f'(a)|^q \right)^{\frac{1}{q}} \right. \\
& \quad \left. + \left( \frac{\alpha(2\alpha^\alpha + (\alpha+1)^\alpha)}{2(\alpha+1)^{\alpha+2}(\alpha+2)} |f'(b)|^q + \frac{4\alpha^{\alpha+1} - \alpha(\alpha+1)^\alpha}{2(\alpha+1)^{\alpha+2}(\alpha+2)} |f'(a)|^q \right)^{\frac{1}{q}} \right].
\end{aligned}$$

This completes the proof.  $\square$

**Remark.** In Theorem 4.5, if one takes  $\alpha = 1$ , one has the following midpoint type inequality

$$\begin{aligned}
& \left| \frac{1}{b-a} \int_a^b f(u) du - f\left(\frac{a+b}{2}\right) \right| \\
& \leq \frac{b-a}{8} \left[ \left( \frac{|f'(b)|^q + 2|f'(a)|^q}{3} \right)^{\frac{1}{q}} + \left( \frac{2|f'(b)|^q + |f'(a)|^q}{3} \right)^{\frac{1}{q}} \right] \quad (4.6)
\end{aligned}$$

**Theorem 4.6.** Let  $a, b \in \mathbb{R}$  with  $a < b$  and  $f : [a, b] \rightarrow \mathbb{R}$  be a differentiable function on  $(a, b)$ . If  $|f'|^q$  is convex on  $[a, b]$  for  $q > 1$ , then the following right Riemann-Liouville fractional integral inequality holds:

$$\begin{aligned}
& \left| \frac{\Gamma(\alpha+1)}{(b-a)^\alpha} J_{b-}^\alpha f(a) - f\left(\frac{a+\alpha b}{\alpha+1}\right) \right| \\
& \leq (b-a) \left[ R_{11}^{\frac{1}{p}}(\alpha, p) \left( \frac{\alpha^2}{2(\alpha+1)^2} |f'(b)|^q + \frac{\alpha^2+2\alpha}{2(\alpha+1)^2} |f'(a)|^q \right)^{\frac{1}{q}} \right. \\
& \quad \left. + R_{12}^{\frac{1}{p}}(\alpha, p) \left( \frac{2\alpha+1}{2(\alpha+1)^2} |f'(b)|^q + \frac{1}{2(\alpha+1)^2} |f'(a)|^q \right)^{\frac{1}{q}} \right] \quad (4.7)
\end{aligned}$$

where

$$\begin{aligned} R_{11}(\alpha, p) &= \int_0^{\frac{\alpha}{\alpha+1}} t^{\alpha p} dt, \\ R_{12}(\alpha, p) &= \int_{\frac{\alpha}{\alpha+1}}^1 (1-t)^p dt, \end{aligned}$$

with  $\frac{1}{p} + \frac{1}{q} = 1$  and  $\alpha > 0$ .

*Proof.* Using Lemma 3.2, Hölder inequality and the convexity of  $|f'|^q$ , we have

$$\begin{aligned} & \left| \frac{\Gamma(\alpha+1)}{(b-a)^\alpha} J_{b^-}^\alpha f(a) - f\left(\frac{a+\alpha b}{\alpha+1}\right) \right| \\ & \leq (b-a) \left[ \int_0^{\frac{\alpha}{\alpha+1}} t^\alpha |f'(tb+(1-t)a)| dt + \int_{\frac{\alpha}{\alpha+1}}^1 (1-t) |f'(tb+(1-t)a)| dt \right] \\ & \leq (b-a) \left[ \left( \int_0^{\frac{\alpha}{\alpha+1}} t^{\alpha p} dt \right)^{\frac{1}{p}} \left( \int_0^{\frac{\alpha}{\alpha+1}} |f'(tb+(1-t)a)|^q dt \right)^{\frac{1}{q}} \right. \\ & \quad \left. + \left( \int_{\frac{\alpha}{\alpha+1}}^1 (1-t)^p dt \right)^{\frac{1}{p}} \left( \int_{\frac{\alpha}{\alpha+1}}^1 |f'(tb+(1-t)a)|^q dt \right)^{\frac{1}{q}} \right] \\ & \leq (b-a) \left[ \left( \int_0^{\frac{\alpha}{\alpha+1}} t^{\alpha p} dt \right)^{\frac{1}{p}} \left( \int_0^{\frac{\alpha}{\alpha+1}} [t|f'(b)|^q + (1-t)|f'(a)|^q] dt \right)^{\frac{1}{q}} \right. \\ & \quad \left. + \left( \int_{\frac{\alpha}{\alpha+1}}^1 (1-t)^p dt \right)^{\frac{1}{p}} \left( \int_{\frac{\alpha}{\alpha+1}}^1 [t|f'(b)|^q + (1-t)|f'(a)|^q] dt \right)^{\frac{1}{q}} \right] \\ & \leq (b-a) \left[ R_{11}^{\frac{1}{p}}(\alpha, p) \left( \frac{\alpha^2}{2(\alpha+1)^2} |f'(b)|^q + \frac{\alpha^2+2\alpha}{2(\alpha+1)^2} |f'(a)|^q \right)^{\frac{1}{q}} \right. \\ & \quad \left. + R_{12}^{\frac{1}{p}}(\alpha, p) \left( \frac{2\alpha+1}{2(\alpha+1)^2} |f'(b)|^q + \frac{1}{2(\alpha+1)^2} |f'(a)|^q \right)^{\frac{1}{q}} \right]. \end{aligned}$$

This completes the proof.  $\square$

**Remark.** In Theorem 4.6, if one takes  $\alpha = 1$ , one has the inequality proved in [4, Theorem 2.3].

## 5. CONCLUSION AND OBSERVATION

We have derived new Hermite-Hadamard type inequalities via convex functions involving only the right Riemann-Liouville fractional integral. Two new fractional integral identities are also obtained. It is expected that idea of this article may attract interested readers

## 6. COMPETING INTERESTS

The authors declare that they have no competing interests.

## REFERENCES

- [1] S. S. Dragomir, R. P. Agarwal, *Two inequalities for differentiable mappings and applications to special means of real numbers and to trapezoidal formula*, Appl. Math. Lett., **11**(5) (1998) 91-95.
- [2] J. Hadamard, *Étude sur les propriétés des fonctions entières et en particulier d'une fonction considérée par Riemann*, J. Math. Pures Appl., **58** (1893) 171-215.
- [3] Ch. Hermite, *Sur deux limites d'une intégrale définie*, Mathesis, **3** (1883) 82-83.
- [4] U. S. Kirmaci, *Inequalities for differentiable mappings and applications to special means of real numbers and to midpoint formula*, Appl. Math. Comp. **147** (2004) 137-146.

- [5] A. A. Kilbas, H. M. Srivastava, J. J. Trujillo, *Theory and applications of fractional differential equations*. Elsevier, Amsterdam (2006).
- [6] C. E. M. Pearce, J. Pecaric, *Inequalities for differentiable mappings with application to special means and quadrature formulae*, Appl. Math. Lett., **13** (2000), 51-55.
- [7] A. W. Roberts, D. E. Varberg, *Convex functions*, Academic Press, New York (1973).
- [8] M.Z. Sarıkaya, E. Set, H. Yıldız and N. Başak, *Hermite-Hadamard's inequalities for fractional integrals and related fractional inequalities*, Math. Comp. Model., **57(9)** (2013) 2403-2407.
- [9] Y. Zhou, *Basic theory of fractional differential equations*, World Scientific, New Jersey (2014).

MEHMET KUNT, DÜNYA KARAPINAR

DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCES, KARADENİZ TECHNICAL UNIVERSITY,  
61080, TRABZON, TURKEY

*Email address:* `mkunt@ktu.edu.tr`; `dunyakarapinar@ktu.edu.tr`

SERCAN TURHAN, İMDAT İŞCAN

DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCES AND ARTS, GİRESUN UNIVERSITY, 28200,  
GİRESUN, TURKEY

*Email address:* `sercan.turhan@giresun.edu.tr`; `imdat.iscan@giresun.edu.tr`