

FURTHER GENERALIZATIONS OF GAMMA, BETA AND RELATED FUNCTIONS

RECEP ŞAHİN, OĞUZ YAĞCI, M. BAKİ YAĞBASAN,
İ. ONUR KIYMAZ*, AYŞEGÜL ÇETİNKAYA

ABSTRACT. In this paper, we present further generalizations of gamma and beta functions by adding extra parameters to their integral representations. Then we introduce new generalizations of hypergeometric and confluent hypergeometric functions by using new beta function. We also obtain many interesting properties such as integral transforms, differentiation formulas, Mellin transforms, transformation formulas, differential and difference relations and a summation formula.

1. INTRODUCTION

Recently, many mathematicians obtained different generalizations of gamma and beta functions by adding some extra parameters to their integral representations. They also examined many properties of these functions. For further reading please see [1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12].

Here, we introduce new generalizations of gamma and beta functions by using an exponential function with four parameter p, q, κ, μ as follows:

$$\Gamma_{p,q}^{(\kappa,\mu)}(x) := \int_0^\infty t^{x-1} e^{\left(-\frac{t^\kappa}{p} - \frac{q}{t^\mu}\right)} dt \quad (1.1)$$

$$(\Re(p) > 0, \Re(q) > 0, \Re(\kappa) > 0, \Re(\mu) > 0),$$

$$B_{p,q}^{(\kappa,\mu)}(x, y) := \int_0^1 t^{x-1} (1-t)^{y-1} e^{\left(-\frac{p}{t^\kappa} - \frac{q}{(1-t)^\mu}\right)} dt \quad (1.2)$$

$$(\Re(p) > 0, \Re(q) > 0, \Re(\kappa) > 0, \Re(\mu) > 0).$$

The special cases of (1.1) are $\Gamma_{1,0}^{(1,1)}(x) = \Gamma(x)$, $\Gamma_{1,p}^{(1,1)}(x) = \Gamma_p(x)$ and (1.2) are $B_{0,0}^{(\kappa,\mu)}(x, y) = B(x, y)$, $B_{p,p}^{(1,1)}(x, y) = B_p(x, y)$, $B_{p,p}^{(m,m)}(x, y) = B_{p:m}(x, y)$, $B_{p,q}^{(1,1)}(x, y) = B_{p,q}(x, y)$ where $\Gamma(x)$, $B(x, y)$ are classical gamma and beta functions (see, [13]), $\Gamma_p(x)$, $B_p(x, y)$ are extended gamma and beta functions defined

2000 *Mathematics Subject Classification.* 33B15, 33C05, 33C15.

Key words and phrases. Hypergeometric function; Gamma and Beta functions; Integral representations; Mellin transforms; Difference operator; Summation formula.

©2018 Ilirias Research Institute, Prishtinë, Kosovë.

Submitted March 4, 2018. Published May 20, 2018.

* Corresponding Author.

Communicated by Senlin Guo.

in [1], $B_{p,m}(x, y)$, $B_{p,q}(x, y)$ are the generalized beta functions given in [5], [3] respectively.

Throughout this paper, we assume that $\min\{\Re(p), \Re(q), \Re(\kappa), \Re(\mu)\} > 0$.

Using $B_{p,q}^{(\kappa,\mu)}(x, y)$, we obtain further generalizations of Gauss and confluent hypergeometric functions

$$F_{p,q}^{(\kappa,\mu)}(\alpha, \beta; \gamma; z) := \sum_{n=0}^{\infty} (\alpha)_n \frac{B_{p,q}^{(\kappa,\mu)}(\beta+n, \gamma-\beta)}{B(\beta, \gamma-\beta)} \frac{z^n}{n!}, \quad |z| < 1 \quad (1.3)$$

$$\Phi_{p,q}^{(\kappa,\mu)}(\beta; \gamma; z) := \sum_{n=0}^{\infty} \frac{B_{p,q}^{(\kappa,\mu)}(\beta+n, \gamma-\beta)}{B(\beta, \gamma-\beta)} \frac{z^n}{n!} \quad (1.4)$$

where $\Re(\gamma) > \Re(\beta) > 0$.

2. INTEGRAL REPRESENTATIONS

Theorem 2.1. *The integral representations of (1.3) and (1.4) are*

$$F_{p,q}^{(\kappa,\mu)}(\alpha, \beta; \gamma; z) = \frac{1}{B(\beta, \gamma-\beta)} \times \int_0^1 t^{\beta-1} (1-t)^{\gamma-\beta-1} e^{\left(-\frac{p}{t^\kappa} - \frac{q}{(1-t)^\mu}\right)} (1-zt)^{-\alpha} dt, \quad (2.1)$$

$$F_{p,q}^{(\kappa,\mu)}(\alpha, \beta; \gamma; z) = \frac{1}{B(\beta, \gamma-\beta)} \times \int_0^\infty u^{\beta-1} (1+u)^{\alpha-\gamma} e^{\left(-\frac{p(u+1)^\kappa}{u^\kappa} - q(u+1)^\mu\right)} (1+u(1-z))^{-\alpha} du, \quad (2.2)$$

$$F_{p,q}^{(\kappa,\mu)}(\alpha, \beta; \gamma; z) = \frac{2}{B(\beta, \gamma-\beta)} \times \int_0^{\frac{\pi}{2}} \frac{\sin^{2\beta-1} \theta \cos^{2\gamma-2\beta-1} \theta}{(1-z \sin^2 \theta)^{-\alpha}} e^{(-p \csc^{2\kappa} \theta - q \sec^{2\mu} \theta)} d\theta, \quad (2.3)$$

$$F_{p,q}^{(\kappa,\mu)}(\alpha, \beta; \gamma; z) = \frac{1}{B(\beta, \gamma-\beta)} \int_0^\infty \left[\sinh^{2\beta-2} \theta \cosh^{2\alpha-2\gamma-2} \theta \times e^{(-p \coth^{2\kappa} \theta - q \cosh^{2\mu} \theta)} (\cosh^2 \theta - z \sinh^2 \theta)^{-\alpha} d\theta \right], \quad (2.4)$$

$$\Phi_{p,q}^{(\kappa,\mu)}(\beta; \gamma; z) = \frac{1}{B(\beta, \gamma-\beta)} \int_0^1 t^{\beta-1} (1-t)^{\gamma-\beta-1} e^{\left(zt - \frac{p}{t^\kappa} - \frac{q}{(1-t)^\mu}\right)} dt, \quad (2.5)$$

$$\Phi_{q,p}^{(\mu,\kappa)}(\beta; \gamma; z) = \frac{e^z}{B(\beta, \gamma-\beta)} \int_0^1 t^{\gamma-\beta-1} (1-t)^{\beta-1} e^{-\left(zt + \frac{p}{(1-t)^\kappa} + \frac{q}{t^\mu}\right)} dt. \quad (2.6)$$

Proof. The integral formulas (2.1) and (2.5) are easily obtained by replacing the definition of generalized beta function (1.2) in (1.3) and (1.4). The integrals in (2.2), (2.3) and (2.4) can be found by making transformations $t = \frac{u}{u+1}$, $t = \sin^2 \theta$, $t = \tanh^2 \theta$ in (2.1), respectively. The integral formula (2.6) is also obtained by changing t with $1-t$ in (2.5). \square

3. DIFFERENTIATION FORMULAS

Theorem 3.1. *The differentiation formulas of (1.3) and (1.4) are*

$$\begin{aligned} \frac{d^n}{dz^n} F_{p,q}^{(\kappa,\mu)}(\alpha, \beta; \gamma; z) &= \frac{(\alpha)_n (\beta)_n}{(\gamma)_n} F_{p,q}^{(\kappa,\mu)}(\alpha + n, \beta + n; \gamma + n; z), \\ \frac{d^n}{dz^n} \Phi_{p,q}^{(\kappa,\mu)}(\beta; \gamma; z) &= \frac{(\beta)_n}{(\gamma)_n} \Phi_{p,q}^{(\kappa,\mu)}(\beta + n; \gamma + n; z). \end{aligned} \quad (3.1)$$

Proof. Differentiation formulas can be found by taking the derivatives of (1.3) and (1.4), using the identities

$$\begin{aligned} B(x, y - x) &= \frac{y}{x} B(x + 1, y - x), \\ (\nu)_{n+1} &= \nu(\nu + 1)_n \end{aligned}$$

which given in [1] and induction. \square

4. MELLIN TRANSFORMS

Theorem 4.1. *The following double Mellin transformation of generalized beta function is*

$$\begin{aligned} \mathfrak{M} \left[B_{p,q}^{(\kappa,\mu)}(x, y); r, s \right] &= \Gamma(s) \Gamma(r) B(x + \kappa s, y + \mu r) \\ &(\Re(s) > 0, \Re(r) > 0, \Re(\beta + r) > 0, \Re(\gamma + s) > 0). \end{aligned}$$

Proof. Double Mellin transform of generalized beta function is

$$\mathfrak{M} \left[B_{p,q}^{(\kappa,\mu)}(x, y); r, s \right] = \int_0^\infty \int_0^\infty q^{r-1} p^{s-1} B_{p,q}^{(\kappa,\mu)}(x, y) dp dq.$$

Considering (1.2) in above equality and making necessary calculations we have

$$\begin{aligned} \mathfrak{M} \left[B_{p,q}^{(\kappa,\mu)}(x, y); r, s \right] &= \int_0^\infty \int_0^\infty q^{r-1} p^{s-1} \\ &\quad \times \left[\int_0^1 t^{x-1} (1-t)^{y-1} e^{\left(-\frac{p}{t^\kappa} - \frac{q}{(1-t)^\mu}\right)} dt \right] dp dq \\ &= \int_0^1 t^{x-1} (1-t)^{y-1} \\ &\quad \times \left[\int_0^\infty q^{r-1} e^{\left(-\frac{q}{(1-t)^\mu}\right)} dq \int_0^\infty p^{s-1} e^{\left(-\frac{p}{t^\kappa}\right)} dp \right] dt \\ &= \Gamma(s) \Gamma(r) \int_0^1 t^{x+\kappa s-1} (1-t)^{y+\mu r-1} dt \\ &= \Gamma(s) \Gamma(r) B(x + \kappa s, y + \mu r). \end{aligned}$$

\square

Theorem 4.2. For $x_1 > 0, x_2 > 0$, the following Mellin-Barnes contour integrals hold true:

$$\begin{aligned} F_{p,q}^{(\kappa,\mu)}(\alpha, \beta; \gamma; z) &= \frac{1}{(2\pi i)^2 B(\beta, \gamma - \beta)} \int_{x_1 - i\infty}^{x_1 + i\infty} \int_{x_2 - i\infty}^{x_2 + i\infty} \left[\Gamma(s) \Gamma(r) \right. \\ &\quad \times B(\beta + \kappa s, \gamma + \kappa s + \mu r) \\ &\quad \left. \times F(\alpha, \beta + \kappa s; \gamma + \kappa s + \mu r; z) q^{-\kappa s} p^{-\mu r} dr ds \right], \end{aligned} \quad (4.1)$$

$$\begin{aligned} \Phi_{p,q}^{(\kappa,\mu)}(\beta; \gamma; z) &= \frac{1}{(2\pi i)^2 B(\beta, \gamma - \beta)} \int_{x_1 - i\infty}^{x_1 + i\infty} \int_{x_2 - i\infty}^{x_2 + i\infty} \left[\Gamma(s) \Gamma(r) \right. \\ &\quad \times B(\beta + \kappa s, \gamma + \kappa s + \mu r) \\ &\quad \left. \times \Phi(\beta + \kappa s; \gamma + \kappa s + \mu r; z) q^{-\kappa s} p^{-\mu r} dr ds \right]. \end{aligned} \quad (4.2)$$

Proof. The double Mellin transform of generalized beta function is

$$\begin{aligned} \mathfrak{M}\left[F_{p,q}^{(\kappa,\mu)}(\alpha, \beta; \gamma; z); r, s\right] &= \int_0^\infty \int_0^\infty p^{s-1} q^{r-1} F_{p,q}^{(\kappa,\mu)}(\alpha, \beta; \gamma; z) dp dq \\ &= \frac{1}{B(\beta, \gamma - \beta)} \int_0^1 \left[t^{\beta-1} (1-t)^{\gamma-\beta-1} (1-zt)^{-\alpha} \right. \\ &\quad \left. \times \int_0^\infty q^{r-1} e^{\left(-\frac{q}{(1-t)^\mu}\right)} dq \int_0^\infty p^{s-1} e^{\left(-\frac{p}{t^\kappa}\right)} dp \right] dt. \end{aligned}$$

Using (2.1) in above equality and making necessary calculations we obtain

$$\begin{aligned} \mathfrak{M}\left[F_{p,q}^{(\kappa,\mu)}(\alpha, \beta; \gamma; z); r, s\right] &= \frac{\Gamma(r) \Gamma(s)}{B(\beta, \gamma - \beta)} \int_0^1 t^{\beta+\kappa s-1} (1-t)^{\gamma-\beta+\mu r-1} (1-zt)^{-\alpha} dt \\ &= \frac{\Gamma(r) \Gamma(s) B(\beta + \kappa s, \gamma + \kappa s + \mu r)}{B(\beta, \gamma - \beta)} F(\alpha, \beta + \kappa s; \gamma + \kappa s + \mu r; z). \end{aligned} \quad (4.3)$$

Using inverse Mellin transforms of (4.3) we have (4.1). We can also prove (4.2) with similar calculations. \square

Theorem 4.3. Difference formula of generalized Gauss hypergeometric function is

$$\begin{aligned} &(\beta - 1)B(\beta - 1, \gamma - \beta + 1) F_{p,q}^{(\kappa,\mu)}(\alpha, \beta - 1; \gamma; z) \\ &= (\gamma - \beta - 1) B(\beta, \gamma - \beta - 1) F_{p,q}^{(\kappa,\mu)}(\alpha, \beta; \gamma - 1; z) \\ &\quad - \alpha z B(\beta, \gamma - \beta) F_{p,q}^{(\kappa,\mu)}(\alpha + 1, \beta; \gamma; z) \\ &\quad - \kappa p B(\beta - \kappa - 1, \gamma - \beta) F_{p,q}^{(\kappa,\mu)}(\alpha, \beta - \kappa - 1; \gamma - \kappa - 1; z) \\ &\quad + \mu q B(\beta, \gamma - \beta - \mu - 1) F_{p,q}^{(\kappa,\mu)}(\alpha, \beta; \gamma - \mu - 1; z). \end{aligned}$$

Proof. We have

$$\begin{aligned} & \frac{d}{dt} \left[H(1-t)(1-t)^{\gamma-\beta-1}(1-zt)^{-\alpha} e^{\left(-\frac{p}{t^\kappa} - \frac{q}{(1-t)^\mu}\right)} \right] \\ &= -(1-zt)^{-\alpha}(1-t)^{\gamma-\beta-1} H(1-t) \left[\delta(1-t) + \frac{(\gamma-\beta-1)}{(1-t)} \right. \\ & \quad \left. - \alpha z(1-zt)^{-1} - \left(\frac{\kappa p}{t^{\kappa+1}} - \frac{\mu q}{(1-t)^{\mu+1}} \right) \right] e^{\left(-\frac{p}{t^\kappa} - \frac{q}{(1-t)^\mu}\right)}. \end{aligned}$$

Taking Mellin transform of both sides and using the following equalities

$$\mathfrak{M}[f'(t); r] = -(r-1)\mathfrak{M}[f(t); r-1]$$

and

$$\begin{aligned} & \mathfrak{M} \left[\frac{d}{dt} \left(H(1-t)(1-t)^{\gamma-\beta-1}(1-zt)^{-\alpha} e^{\left(-\frac{p}{t^\kappa} - \frac{q}{(1-t)^\mu}\right)} \right); \beta \right] \\ &= B(\beta, \gamma - \beta) F_{p,q}^{(\kappa, \mu)}(\alpha, \beta; \gamma; z) \end{aligned}$$

in [13], we get the desired result. \square

The next theorem can be proved with similar calculations of the above theorem. That's why we omit its proof.

Theorem 4.4. *Difference formula of generalized confluent hypergeometric function is*

$$\begin{aligned} & (\beta-1)B(\beta-1, \gamma-\beta+1) \Phi_{p,q}^{(\kappa, \mu)}(\beta-1; \gamma; z) = -zB(\beta, \gamma-\beta) \Phi_{p,q}^{(\kappa, \mu)}(\beta; \gamma; z) \\ & + (\gamma-\beta-1)B(\beta, \gamma-\beta-1) \Phi_{p,q}^{(\kappa, \mu)}(\beta; \gamma-1; z) \\ & - \kappa p B(\beta-\kappa-1, \gamma-\beta) \Phi_{p,q}^{(\kappa, \mu)}(\beta-\kappa-1; \gamma-\kappa-1; z) \\ & + \mu q B(\beta, \gamma-\beta-\mu-1) \Phi_{p,q}^{(\kappa, \mu)}(\beta; \gamma-\mu-1; z). \end{aligned}$$

5. TRANSFORMATION FORMULAS

Theorem 5.1. *The transformation formulas of (1.3) and (1.4) are*

$$F_{p,q}^{(\kappa, \mu)}(\alpha, \beta; \gamma; z) = (1-z)^{-\alpha} F_{q,p}^{(\mu, \kappa)}\left(\alpha, \gamma-\beta; \gamma; -\frac{z}{1-z}\right) \quad (5.1)$$

$$(|\arg(1-z)| < \pi, \Re(\gamma) > \Re(\beta) > 0),$$

$$F_{p,q}^{(\kappa, \mu)}\left(\alpha, \beta; \gamma; 1-\frac{1}{z}\right) = z^\alpha F_{q,p}^{(\mu, \kappa)}(\alpha, \gamma-\beta; \gamma; 1-z) \quad (5.2)$$

$$(|\arg(1-z)| < \pi, \Re(\gamma) > \Re(\beta) > 0),$$

$$F_{p,q}^{(\kappa, \mu)}\left(\alpha, \beta; \gamma; \frac{z}{z+1}\right) = (1+z)^\alpha F_{q,p}^{(\mu, \kappa)}(\alpha, \gamma-\beta; \gamma; -z) \quad (5.3)$$

$$(|\arg(1-z)| < \pi, \Re(\gamma) > \Re(\beta) > 0),$$

$$\Phi_{p,q}^{(\kappa, \mu)}(\beta; \gamma; z) = e^z \Phi_{q,p}^{(\mu, \kappa)}(\gamma-\beta; \gamma; -z) \quad (5.4)$$

$$(\Re(\gamma) > \Re(\beta) > 0).$$

Proof. By using the equality

$$[1 - z(1 - u)]^{-\alpha} = (1 - z)^{-\alpha} \left(1 + \frac{z}{1 - z}u\right)^{-\alpha}$$

in (2.1) and taking $1 - u$ in stead of t , we find

$$F_{q,p}^{(\mu,\kappa)}(\alpha, \beta; \gamma; z) = \frac{(1 - z)^{-\alpha}}{B(\beta, \gamma - \beta)} \int_0^1 \left[u^{\gamma - \beta - 1} (1 - u)^{\beta - 1} \times e^{\left(-\frac{q}{u^\mu} - \frac{p}{(1-u)^\kappa}\right)} \left(1 + \frac{z}{1 - z}u\right)^{-\alpha} du \right].$$

This proves (5.1). For (5.2) and (5.3) one can replace z by $1 - \frac{1}{z}$ and $\frac{z}{z+1}$ in (5.1), respectively. Finally, equation (5.4) can be easily obtained from (2.5) and (2.6). \square

6. DIFFERENTIAL AND DIFFERENCE RELATIONS

Theorem 6.1. *The differential and difference relations of (1.3) and (1.4) are*

$$\Delta_\alpha F_{p,q}^{(\kappa,\mu)}(\alpha, \beta; \gamma; z) = \frac{bz}{c} F_{p,q}^{(\kappa,\mu)}(\alpha + 1, \beta + 1; \gamma + 1; z), \quad (6.1)$$

$$\frac{d}{dz} F_{p,q}^{(\kappa,\mu)}(\alpha, \beta; \gamma; z) = \frac{\alpha}{z} \Delta_\alpha F_{p,q}^{(\kappa,\mu)}(\alpha, \beta; \gamma; z), \quad (6.2)$$

$$\beta \Delta_\beta \Phi_{p,q}^{(\kappa,\mu)}(\beta; \gamma + 1; z) + \gamma \Delta_\gamma \Phi_{p,q}^{(\kappa,\mu)}(\beta; \gamma; z) = 0, \quad (6.3)$$

$$\frac{d}{dz} \Phi_{p,q}^{(\kappa,\mu)}(\beta; \gamma; z) = \frac{\beta}{\gamma} \Phi_{p,q}^{(\kappa,\mu)}(\beta; \gamma + 1; z) - \Delta_\gamma \Phi_{p,q}^{(\kappa,\mu)}(\beta; \gamma; z), \quad (6.4)$$

where Δ_α is the well-known difference operator

$$\Delta_\alpha f(\alpha, \dots) = f(\alpha + 1, \dots) - f(\alpha, \dots).$$

Proof. In (2.1), if we write $\alpha + 1$, $\beta + 1$, and $\gamma + 1$ in stead of α , β , and γ and using

$$\Delta_\alpha F_{p,q}^{(\kappa,\mu)}(\alpha, \beta; \gamma; z) = F_{p,q}^{(\kappa,\mu)}(\alpha + 1, \beta; \gamma; z) - F_{p,q}^{(\kappa,\mu)}(\alpha, \beta; \gamma; z)$$

we obtain (6.1). To prove (6.2) we can use (3.1) in (6.1). By making similar calculations we can prove (6.3) and (6.4). \square

7. SUMMATION FORMULA

We also give a summation formula for the special case of generalized Gauss hypergeometric function when $z = 1$.

Theorem 7.1. *The summation formula of (1.3) is*

$$F_{p,q}^{(\kappa,\mu)}(\alpha, \beta; \gamma; 1) = \frac{B_{p,q}^{(\kappa,\mu)}(\beta, \gamma - \alpha - \beta)}{B(\beta, \gamma - \beta)} \\ (\Re(\gamma - \beta - \alpha) > 0).$$

Proof. The result can obtained easily from (2.1) by taking $z = 1$ and using (1.2). \square

Acknowledgements: The authors are grateful to the anonymous referees for their careful reading that led to a better presentation of the paper.

REFERENCES

- [1] M. A. Chaudhry, A. Qadir, M. Raque and S. M. Zubair, *Extension of Euler's Beta function*, J. Comput. Appl. Math. **78** (1997) 19–32.
- [2] M. A. Chaudhry, A. Qadir, H. M. Srivastava and R. B. Paris, *Extended hypergeometric and confluent hypergeometric functions*, Appl. Math. Comput. **159** (2004) 589–602.
- [3] J. Choi, A. K. Rathie and R. K. Parmar, *Extension of extended beta, hypergeometric and confluent hypergeometric functions*, Honam Mathematical Journal **36** (2) (2014) 357–385.
- [4] A. Çetinkaya, İ. O. Kıymaz, P. Agarwal and R. Agarwal, *A comparative study on generating function relations for generalized hypergeometric functions via generalized fractional operators*, Advances in Difference Equations **2018:156** (2018) <https://doi.org/10.1186/s13662-018-1612-0>.
- [5] Dong-Myung Lee, A. K. Rathie, R. K. Parmar and Y. S. Kim, *Generalization of extended beta function, hypergeometric and confluent hypergeometric functions*, Honam Mathematical Journal **33** (2) (2011) 187–206.
- [6] M. J. Luo, G. V. Milovanovic and P. Agarwal, *Some results on the extended beta and extended hypergeometric functions*, Appl. Math. Comput. **248** (2014) 631–651.
- [7] E. Özergin, *Some Properties of Hypergeometric Functions*, Ph.D. Thesis, Eastern Mediterranean University, North Cyprus, Turkey, (2011).
- [8] E. Özergin, M. A. Özarlan and A. Altın, *Extension of Gamma, beta and hypergeometric functions*, Journal of Comp. and Applied Math. **235** (2011) 4601–4610.
- [9] R. K. Parmar, *A new generalization of Gamma, Beta, hypergeometric and Confluent hypergeometric functions*, Le Matematiche **LXVIII** (2013) 33–52.
- [10] H. M. Srivastava, P. Agarwal and S. Jain, *Generating functions for the generalized Gauss hypergeometric functions*, Appl. Math. Comput. **247** (2014) 348–352.
- [11] H. M. Srivastava, A. Çetinkaya and İ. O. Kıymaz, *A Certain generalized Pochhammer symbol and its applications to hypergeometric functions*, Appl. Math. Comput. **226** (2014) 484–491.
- [12] H. M. Srivastava, R. Jain and M. K. Bansal, *A Study of the S-Generalized Gauss Hypergeometric Function and Its Associated Integral Transforms*, Turkish Journal of Analysis and Number Theory **3** (5) (2015) 116–119.
- [13] H. M. Srivastava and P. W. Karlsson, *Multiple Gaussian Hypergeometric Series*, Halsted Press (Ellis Horwood Limited, Chichester), John Wiley and Sons, New York, Chichester, Brisbane and Toronto, (1985).

R. ŞAHİN

KIRIKKALE UNIVERSITY, DEPT. OF MATHEMATICS, KIRIKKALE-TURKEY
E-mail address: receptsahin@kku.edu.tr

O. YAĞCI

KIRIKKALE UNIVERSITY, DEPT. OF MATHEMATICS, KIRIKKALE-TURKEY
E-mail address: oguzyagci26@gmail.com

M. B. YAĞBASAN

AHI EVRAN UNIV., DEPT. OF MATHEMATICS, KIRŞEHİR-TURKEY
E-mail address: mbyagbasan@ahievran.edu.tr

İ. O. KIYMAZ

AHI EVRAN UNIV., DEPT. OF MATHEMATICS, KIRŞEHİR-TURKEY
E-mail address: iokiyamaz@ahievran.edu.tr

A. ÇETİNKAYA

AHI EVRAN UNIV., DEPT. OF MATHEMATICS, KIRŞEHİR-TURKEY
E-mail address: acetinkaya@ahievran.edu.tr