

**q -DEFORMED AND λ -PARAMETRIZED A -GENERALIZED
LOGISTIC FUNCTION BASED COMPLEX VALUED
TRIGONOMETRIC AND HYPERBOLIC NEURAL NETWORK
HIGH ORDER APPROXIMATIONS**

GEORGE A. ANASTASSIOU

ABSTRACT. Here we research the univariate quantitative approximation of complex valued continuous functions on a compact interval by complex valued neural network operators. These approximations are derived by establishing Jackson type inequalities involving the modulus of continuity of the engaged function's high order derivatives. The nature of our approximations are trigonometric and hyperbolic. Our operators are defined by using a density function generated by a q -deformed and λ -parametrized A -generalized logistic function, which is a sigmoid function. The approximations are pointwise and of the uniform norm. The related complex valued feed-forward neural networks are with one hidden layer.

1. INTRODUCTION

The author in [1] and [2], see Chapters 2-5, was the first to establish neural network approximation to continuous functions with rates by very specifically defined neural network operators of Cardaliaguet-Euvrard and "Squashing" types, by employing the modulus of continuity of the engaged function or its high order derivative, and producing very tight Jackson type inequalities. He treats there both the univariate and multivariate cases. The defining these operators "bell-shaped" and "squashing" functions are assumed to be of compact support.

Again the author inspired by [12], continued his studies on neural networks approximation by introducing and using the proper quasi-interpolation operators of sigmoidal and hyperbolic tangent type which resulted into [3]-[7], by treating both the univariate and multivariate cases.

Let h be a general sigmoid activation function with $h(0) = 0$, and $y = \pm 1$ the horizontal asymptotes. Of course h is strictly increasing over \mathbb{R} . Let the parameter $0 < r < 1$ and $x > 0$. Then clearly $-x < x$ and $-x < -rx < rx < x$, furthermore

1991 *Mathematics Subject Classification.* 41A17, 41A25, 41A99, 42A10.

Key words and phrases. q -deformed and λ -parametrized A -generalized logistic function, complex valued neural network approximation, complex valued quasi-interpolation operator, modulus of continuity, trigonometric and hyperbolic high order approximation.

©2023 Ilirias Research Institute, Prishtinë, Kosovë.

Submitted June 7, 2023. Published June 20, 2023.

Communicated by M.A.Noor.

it holds $h(-x) < h(-rx) < h(rx) < h(x)$. Consequently the sigmoid $y = h(rx)$ has a graph inside the graph of $y = h(x)$, of course with the same asymptotes $y = \pm 1$. Therefore $h(rx)$ has derivatives (gradients) at more points x than $h(x)$ has different than zero or not as close to zero, thus killing less number of neurons! And of course $h(rx)$ is more distant from $y = \pm 1$, than $h(x)$ it is. A highly desired fact in Neural Networks theory.

Different activation functions allow for different non-linearities which might work better for solving a specific function. So the need to use neural networks with various activation functions is vivid. Thus, performing neural network approximations using different activation functions is not only necessary but fully justified.

Also brain asymmetry has been observed in animals and humans in terms of structure, function and behaviour. This lateralization is thought to reflect evolutionary, hereditary, developmental, experiential and pathological factors. Therefore it is natural to consider for our study deformed neural network activation functions and operators. So this paper is a specific study under this philosophy of approaching reality as close as possible.

Consequently the author here performs q -deformed and λ -parametrized A -generalized logistic function activated high order neural network approximations to continuous functions over compact intervals of the real line with complex values. All convergences are with rates expressed via the modulus of continuity of the involved functions high order derivatives, deriving by very tight Jackson type inequalities.

The basis of our higher order approximations here are some newly discovered by the author trigonometric and hyperbolic type Taylor's formulae.

Our compact intervals are not necessarily symmetric to the origin. In preparation to prove our results we describe important properties of the basic density function defining our operators which is induced by a q -deformed and λ -parametrized A -generalized logistic function, which is a sigmoid activation function.

Feed-forward X -valued neural networks (FNNs) with one hidden layer, the only type of networks we deal with in this article, are mathematically expressed as

$$N_n(x) = \sum_{j=0}^n c_j \sigma(\langle a_j \cdot x \rangle + b_j), \quad x \in \mathbb{R}^s, \quad s \in \mathbb{N},$$

where for $0 \leq j \leq n$, $b_j \in \mathbb{R}$ are the thresholds, $a_j \in \mathbb{R}^s$ are the connection weights, $c_j \in X$ are the coefficients, $\langle a_j \cdot x \rangle$ is the inner product of a_j and x , and σ is the activation function of the network. About neural networks in general read [13], [15], [16].

2. PRELIMINARIES

The following background comes from [10].

We consider here the q -deformed and λ -parametrized function

$$\varphi_{q,\lambda}(x) = \frac{1}{1 + qA^{-\lambda x}}, \quad x \in \mathbb{R}, \quad \text{where } q, \lambda > 0, \quad A > 1. \quad (1)$$

This is an A -generalized logistic type function.

We easily observe that

$$\varphi_{q,\lambda}(+\infty) = 1, \quad \varphi_{q,\lambda}(-\infty) = 0. \quad (2)$$

Furthermore we have

$$1 - \varphi_{\frac{1}{q},\lambda}(-x) = 1 - \frac{1}{1 + \frac{1}{q}A^{\lambda x}} = \frac{1 + \frac{1}{q}A^{\lambda x} - 1}{1 + \frac{1}{q}A^{\lambda x}} = \frac{\frac{1}{q}A^{\lambda x}}{1 + \frac{1}{q}A^{\lambda x}} = \frac{1}{\frac{1}{\frac{1}{q}A^{\lambda x}} + 1} = \frac{1}{1 + qA^{-\lambda x}} = \varphi_{q,\lambda}(x),$$

proving

$$\varphi_{q,\lambda}(x) = 1 - \varphi_{\frac{1}{q},\lambda}(-x). \quad (3)$$

We also have that

$$\varphi_{q,\lambda}(0) = \frac{1}{1 + q}. \quad (4)$$

Consider the activation function

$$G_{q,\lambda}(x) := \frac{1}{2}(\varphi_{q,\lambda}(x+1) - \varphi_{q,\lambda}(x-1)), \quad x \in \mathbb{R}. \quad (5)$$

Then

$$\begin{aligned} G_{q,\lambda}(-x) &= \frac{1}{2}(\varphi_{q,\lambda}(-x+1) - \varphi_{q,\lambda}(-x-1)) = \\ &= \frac{1}{2}\left(1 - \varphi_{\frac{1}{q},\lambda}(x-1) - 1 + \varphi_{\frac{1}{q},\lambda}(x+1)\right) = \\ &= \frac{1}{2}\left(\varphi_{\frac{1}{q},\lambda}(x+1) - \varphi_{\frac{1}{q},\lambda}(x-1)\right) = G_{\frac{1}{q},\lambda}(x). \end{aligned} \quad (6)$$

That is

$$G_{q,\lambda}(-x) = G_{\frac{1}{q},\lambda}(x), \quad \forall x \in \mathbb{R}. \quad (7)$$

We have

$$\begin{aligned} \varphi'_{q,\lambda}(x) &= \left((1 + qA^{-\lambda x})^{-1}\right)' = \\ &= -1(1 + qA^{-\lambda x})^{-2} q(\ln A) A^{-\lambda x} (-\lambda) = q\lambda(\ln A)(1 + qA^{-\lambda x})^{-2} A^{-\lambda x} > 0. \end{aligned} \quad (8)$$

So that $\varphi_{q,\lambda}$ is a strictly increasing function over \mathbb{R} .

Hence it holds

$$\begin{aligned} \varphi'_{q,\lambda}(x) &= \frac{q\lambda(\ln A)}{(1 + qA^{-\lambda x})^2 A^{\lambda x}} = \\ &= \frac{q\lambda(\ln A)}{(1 + q^2 A^{-2\lambda x} + 2qA^{-\lambda x}) A^{\lambda x}} = \frac{q\lambda(\ln A)}{(A^{\lambda x} + q^2 A^{-\lambda x} + 2q)}. \end{aligned} \quad (9)$$

That is

$$\varphi'_{q,\lambda}(x) = q\lambda(\ln A)(A^{\lambda x} + q^2 A^{-\lambda x} + 2q)^{-1}. \quad (10)$$

Therefore it holds

$$\begin{aligned} \varphi''_{q,\lambda}(x) &= q\lambda(\ln A)(-1)(A^{\lambda x} + q^2 A^{-\lambda x} + 2q)^{-2} ((\ln A) A^{\lambda x} \lambda + q^2 (\ln A) A^{-\lambda x} (-\lambda)) = \\ &= q\lambda^2 (\ln A)^2 (A^{\lambda x} + q^2 A^{-\lambda x} + 2q)^{-2} (q^2 A^{-\lambda x} - A^{\lambda x}). \end{aligned} \quad (11)$$

That is

$$\varphi''_{q,\lambda}(x) = q\lambda^2 (\ln A)^2 (A^{\lambda x} + q^2 A^{-\lambda x} + 2q)^{-2} (q^2 A^{-\lambda x} - A^{\lambda x}) \in C(\mathbb{R}). \quad (12)$$

We have

$$\begin{aligned} q^2 A^{-\lambda x} - A^{\lambda x} > 0, & \text{ iff } q^2 A^{-\lambda x} > A^{\lambda x}, \text{ iff } q^2 > A^{2\lambda x}, \text{ iff } q > A^{\lambda x}, \\ & \text{ iff } \log_A q > \lambda x, \text{ iff } x < \frac{\log_A q}{\lambda}. \end{aligned}$$

So, $\varphi''_{q,\lambda}(x) > 0$, for $x < \frac{\log_A q}{\lambda}$ and there $\varphi_{q,\lambda}$ is concave up.

When $x > \frac{\log_A q}{\lambda}$, we have $\varphi''_{q,\lambda}(x) < 0$ and $\varphi_{q,\lambda}$ is concave down.

Of course

$$\varphi''_{q,\lambda}\left(\frac{\log_A q}{\lambda}\right) = 0.$$

So, $\varphi_{q,\lambda}$ is a sigmoid function, see [9].

We have that

$$G'_{q,\lambda}(x) = \frac{1}{2} (\varphi'_{q,\lambda}(x+1) - \varphi'_{q,\lambda}(x-1)).$$

We got that $\varphi'_{q,\lambda}$ is strictly increasing for $x < \frac{\log_A q}{\lambda}$. Let $x < \frac{\log_A q}{\lambda} - 1$, then $x-1 < x+1 < \frac{\log_A q}{\lambda}$. Hence $\varphi'_{q,\lambda}(x+1) > \varphi'_{q,\lambda}(x-1)$. Thus $G'_{q,\lambda} > 0$, i.e. $G_{q,\lambda}$ is strictly increasing over $(-\infty, \frac{\log_A q}{\lambda} - 1)$.

Let now $x > \frac{\log_A q}{\lambda} + 1$, then $x+1 > x-1 > \frac{\log_A q}{\lambda}$, and $\varphi'_{q,\lambda}(x+1) < \varphi'_{q,\lambda}(x-1)$, by $\varphi'_{q,\lambda}$ being strictly decreasing over $(\frac{\log_A q}{\lambda}, +\infty)$. Hence $G'_{q,\lambda} < 0$, and $G_{q,\lambda}$ is strictly decreasing over $(\frac{\log_A q}{\lambda}, +\infty)$.

Let now $\frac{\log_A q}{\lambda} - 1 \leq x \leq \frac{\log_A q}{\lambda} + 1$. We have that

$$\begin{aligned} G''_{q,\lambda}(x) &= \frac{1}{2} (\varphi''_{q,\lambda}(x+1) - \varphi''_{q,\lambda}(x-1)) = \\ &= \frac{q\lambda^2 (\ln A)^2}{2} \left[\frac{(q^2 A^{-\lambda(x+1)} - A^{\lambda(x+1)})}{(A^{\lambda(x+1)} + q^2 A^{-\lambda(x+1)} + 2q)^2} - \frac{(q^2 A^{-\lambda(x-1)} - A^{\lambda(x-1)})}{(A^{\lambda(x-1)} + q^2 A^{-\lambda(x-1)} + 2q)^2} \right] = \\ &= \frac{q\lambda^2 (\ln A)^2}{2} \left[\frac{(q^2 - A^{2\lambda(x+1)})}{(A^{\lambda(x+1)} + q^2 A^{-\lambda(x+1)} + 2q)^2 A^{\lambda(x+1)}} - \frac{(q^2 - A^{2\lambda(x-1)})}{(A^{\lambda(x-1)} + q^2 A^{-\lambda(x-1)} + 2q)^2 A^{\lambda(x-1)}} \right] = \\ &= \frac{q\lambda^2 (\ln A)^2}{2} \left[\frac{(q - A^{\lambda(x+1)})(q + A^{\lambda(x+1)})}{(A^{\lambda(x+1)} + q^2 A^{-\lambda(x+1)} + 2q)^2 A^{\lambda(x+1)}} - \frac{(q - A^{\lambda(x-1)})(q + A^{\lambda(x-1)})}{(A^{\lambda(x-1)} + q^2 A^{-\lambda(x-1)} + 2q)^2 A^{\lambda(x-1)}} \right]. \end{aligned} \tag{13}$$

By $\frac{\log_A q}{\lambda} \leq x+1 \Leftrightarrow \log_A q \leq \lambda(x+1) \Leftrightarrow q \leq A^{\lambda(x+1)} \Leftrightarrow q - A^{\lambda(x+1)} \leq 0$.

By $x \leq \frac{\log_A q}{\lambda} + 1 \Leftrightarrow x-1 \leq \frac{\log_A q}{\lambda} \Leftrightarrow \lambda(x-1) \leq \log_A q \Leftrightarrow A^{\lambda(x-1)} \leq q \Leftrightarrow q - A^{\lambda(x-1)} \geq 0$.

Clearly, when $\frac{\log_A q}{\lambda} - 1 \leq x \leq \frac{\log_A q}{\lambda} + 1$ by the above we get that $G''_{q,\lambda}(x) \leq 0$, that is $G''_{q,\lambda}$ is concave down there.

Clearly $G_{q,\lambda}$ is strictly concave down over $(\frac{\log_A q}{\lambda} - 1, \frac{\log_A q}{\lambda} + 1)$.

Overall $G_{q,\lambda}$ is a bell-shaped function over \mathbb{R} .

Of course it holds $G''_{q,\lambda}\left(\frac{\log_A q}{\lambda}\right) < 0$.

We have that

$$\begin{aligned} G'_{q,\lambda}\left(\frac{\log_A q}{\lambda}\right) &= \frac{1}{2} \left(\varphi'_{q,\lambda}\left(\frac{\log_A q}{\lambda} + 1\right) - \varphi'_{q,\lambda}\left(\frac{\log_A q}{\lambda} - 1\right) \right) = \\ &= \frac{q\lambda (\ln A)}{2} \left[\frac{1}{A^{\lambda\left(\frac{\log_A q}{\lambda} + 1\right)} + q^2 A^{-\lambda\left(\frac{\log_A q}{\lambda} + 1\right)} + 2q} - \frac{1}{A^{\lambda\left(\frac{\log_A q}{\lambda} - 1\right)} + q^2 A^{-\lambda\left(\frac{\log_A q}{\lambda} - 1\right)} + 2q} \right] = \end{aligned} \tag{14}$$

$$\frac{q\lambda(\ln A)}{2} \left[\frac{A^\lambda \left(\frac{\log_A q}{\lambda} - 1\right) + q^2 A^{-\lambda} \left(\frac{\log_A q}{\lambda} - 1\right) - A^\lambda \left(\frac{\log_A q}{\lambda} + 1\right) - q^2 A^{-\lambda} \left(\frac{\log_A q}{\lambda} + 1\right)}{\left(A^\lambda \left(\frac{\log_A q}{\lambda} + 1\right) + q^2 A^{-\lambda} \left(\frac{\log_A q}{\lambda} + 1\right) + 2q\right) \left(A^\lambda \left(\frac{\log_A q}{\lambda} - 1\right) + q^2 A^{-\lambda} \left(\frac{\log_A q}{\lambda} - 1\right) + 2q\right)} \right] = \quad (15)$$

$$\begin{aligned} & \frac{q\lambda(\ln A)}{2} \left[\frac{qA^{-\lambda} + q^2 q^{-1} A^\lambda - qA^\lambda - q^2 q^{-1} A^{-\lambda}}{(qA^\lambda + q^2 q^{-1} A^{-\lambda} + 2q)(qA^\lambda + q^2 q^{-1} A^{-\lambda} + 2q)} \right] = \\ & \frac{q\lambda(\ln A)}{2} \left[\frac{qA^{-\lambda} + qA^\lambda - qA^\lambda - qA^{-\lambda}}{(qA^\lambda + qA^{-\lambda} + 2q)(qA^\lambda + qA^{-\lambda} + 2q)} \right] = 0. \end{aligned} \quad (16)$$

So $\frac{\log_A q}{\lambda}$ is the only critical number of $G_{q,\lambda}$ over \mathbb{R} . Therefore $G_{q,\lambda} \left(\frac{\log_A q}{\lambda} \right)$ is the maximum of $G_{q,\lambda}$.

We calculate it:

We have that

$$\begin{aligned} G_{q,\lambda} \left(\frac{\log_A q}{\lambda} \right) &= \frac{1}{2} \left(\varphi_{q,\lambda} \left(\frac{\log_A q}{\lambda} + 1 \right) - \varphi_{q,\lambda} \left(\frac{\log_A q}{\lambda} - 1 \right) \right) = \\ & \frac{1}{2} \left(\frac{1}{1 + qA^{-\lambda} \left(\frac{\log_A q}{\lambda} + 1\right)} - \frac{1}{1 + qA^{-\lambda} \left(\frac{\log_A q}{\lambda} - 1\right)} \right) = \quad (17) \\ & \frac{1}{2} \left(\frac{1}{1 + qq^{-1} A^{-\lambda}} - \frac{1}{1 + qq^{-1} A^\lambda} \right) = \frac{1}{2} \left(\frac{1}{1 + A^{-\lambda}} - \frac{1}{1 + A^\lambda} \right) = \\ & \frac{1}{2} \left(\frac{A^\lambda - A^{-\lambda}}{(1 + A^{-\lambda})(1 + A^\lambda)} \right) = \frac{A^\lambda - 1}{2(A^\lambda + 1)}. \end{aligned}$$

The global maximum of $G_{q,\lambda}$ is

$$G_{q,\lambda} \left(\frac{\log_A q}{\lambda} \right) = \frac{A^\lambda - 1}{2(A^\lambda + 1)}. \quad (18)$$

Finally we have that

$$\lim_{x \rightarrow +\infty} G_{q,\lambda}(x) = \frac{1}{2} (\varphi_{q,\lambda}(+\infty) - \varphi_{q,\lambda}(+\infty)) = 0, \quad (19)$$

and

$$\lim_{x \rightarrow -\infty} G_{q,\lambda}(x) = \frac{1}{2} (\varphi_{q,\lambda}(-\infty) - \varphi_{q,\lambda}(-\infty)) = 0. \quad (20)$$

Consequently the x -axis is the horizontal asymptote of $G_{q,\lambda}$. Of course $G_{q,\lambda}(x) > 0$, $\forall x \in \mathbb{R}$.

We need

Theorem 2.1. ([10]) *It holds*

$$\sum_{i=-\infty}^{\infty} G_{q,\lambda}(x-i) = 1, \quad \forall x \in \mathbb{R}, \forall q, \lambda > 0, A > 1. \quad (21)$$

Thus

$$\sum_{i=-\infty}^{\infty} G_{q,\lambda}(nx-i) = 1, \quad \forall n \in \mathbb{N}, \forall x \in \mathbb{R}. \quad (22)$$

Similarly, it holds

$$\sum_{i=-\infty}^{\infty} G_{\frac{1}{q},\lambda}(x-i) = 1, \quad \forall x \in \mathbb{R}. \quad (23)$$

But $G_{\frac{1}{q}, \lambda}(x-i) \stackrel{(7)}{=} G_{q, \lambda}(i-x), \forall x \in \mathbb{R}$.
Hence

$$\sum_{i=-\infty}^{\infty} G_{q, \lambda}(i-x) = 1, \forall x \in \mathbb{R}, \quad (24)$$

and

$$\sum_{i=-\infty}^{\infty} G_{q, \lambda}(i+x) = 1, \forall x \in \mathbb{R}. \quad (25)$$

It follows

Theorem 2.2. ([10]) *It holds*

$$\int_{-\infty}^{\infty} G_{q, \lambda}(x) dx = 1, \quad \lambda, q > 0, A > 1. \quad (26)$$

So that $G_{q, \lambda}$ is a density function on \mathbb{R} ; $\lambda, q > 0, A > 1$.

We need the following result

Theorem 2.3. ([10]) *Let $0 < \alpha < 1$, and $n \in \mathbb{N}$ with $n^{1-\alpha} > 2$. Then*

$$\begin{cases} \sum_{k=-\infty}^{\infty} G_{q, \lambda}(nx-k) < \max\left\{q, \frac{1}{q}\right\} \frac{1}{A^{\lambda(n^{1-\alpha}-2)}} = \gamma A^{-\lambda(n^{1-\alpha}-2)}, \\ : |nx-k| \geq n^{1-\alpha} \end{cases} \quad (27)$$

where $q, \lambda > 0, A > 1; \gamma := \max\left\{q, \frac{1}{q}\right\}$.

Let $\lceil \cdot \rceil$ the ceiling of the number, and $\lfloor \cdot \rfloor$ the integral part of the number.

We also need

Theorem 2.4. ([10]) *Let $x \in [a, b] \subset \mathbb{R}$ and $n \in \mathbb{N}$ so that $\lceil na \rceil \leq \lfloor nb \rfloor$. For $q > 0, \lambda > 0, A > 1$, we consider the number $\lambda_q > z_0 > 0$ with $G_{q, \lambda}(z_0) = G_{q, \lambda}(0)$ and $\lambda_q > 1$. Then*

$$\frac{1}{\sum_{k=\lceil na \rceil}^{\lfloor nb \rfloor} G_{q, \lambda}(nx-k)} < \max\left\{\frac{1}{G_{q, \lambda}(\lambda_q)}, \frac{1}{G_{\frac{1}{q}, \lambda}\left(\frac{\lambda_q}{q}\right)}\right\} =: K(q). \quad (28)$$

We finally mention

Remark 1. ([10]) (i) *We have that*

$$\lim_{n \rightarrow +\infty} \sum_{k=\lceil na \rceil}^{\lfloor nb \rfloor} G_{q, \lambda}(nx-k) \neq 1, \quad \text{for at least some } x \in [a, b], \quad (29)$$

where $\lambda, q > 0$.

(ii) *Let $[a, b] \subset \mathbb{R}$. For large n we always have $\lceil na \rceil \leq \lfloor nb \rfloor$. Also $a \leq \frac{k}{n} \leq b$, iff $\lceil na \rceil \leq k \leq \lfloor nb \rfloor$. In general it holds*

$$\sum_{k=\lceil na \rceil}^{\lfloor nb \rfloor} G_{q, \lambda}(nx-k) \leq 1. \quad (30)$$

Let $(\mathbb{C}, |\cdot|)$ be the Banach space of the complex numbers over the reals.

Definition 1. Let $f \in C([a, b], \mathbb{C})$ and $n \in \mathbb{N} : \lceil na \rceil \leq \lfloor nb \rfloor$. We introduce and define the \mathbb{C} -valued linear neural network operators

$$L_n(f, x) := \frac{\sum_{k=\lceil na \rceil}^{\lfloor nb \rfloor} f\left(\frac{k}{n}\right) G_{q,\lambda}(nx - k)}{\sum_{k=\lceil na \rceil}^{\lfloor nb \rfloor} G_{q,\lambda}(nx - k)}, \quad x \in [a, b]. \quad (31)$$

Clearly here $L_n(f, x) \in C([a, b], \mathbb{C})$.

For convenience we use the same L_n for real valued functions when needed. We study here the high order pointwise and uniform convergence of $L_n(f, x)$ to $f(x)$ with rates.

For convenience, also we call

$$L_n^*(f, x) := \sum_{k=\lceil na \rceil}^{\lfloor nb \rfloor} f\left(\frac{k}{n}\right) G_{q,\lambda}(nx - k), \quad (32)$$

(similarly, L_n^* can be defined for real valued functions) that is

$$L_n(f, x) := \frac{L_n^*(f, x)}{\sum_{k=\lceil na \rceil}^{\lfloor nb \rfloor} G_{q,\lambda}(nx - k)}. \quad (33)$$

So that

$$L_n(f, x) - f(x) = \frac{L_n^*(f, x)}{\sum_{k=\lceil na \rceil}^{\lfloor nb \rfloor} G_{q,\lambda}(nx - k)} - f(x) = \quad (34)$$

$$\frac{L_n^*(f, x) - f(x) \left(\sum_{k=\lceil na \rceil}^{\lfloor nb \rfloor} G_{q,\lambda}(nx - k) \right)}{\sum_{k=\lceil na \rceil}^{\lfloor nb \rfloor} G_{q,\lambda}(nx - k)}.$$

Consequently, we derive that

$$|L_n(f, x) - f(x)| \leq K(q) \left| L_n^*(f, x) - f(x) \left(\sum_{k=\lceil na \rceil}^{\lfloor nb \rfloor} G_{q,\lambda}(nx - k) \right) \right| =$$

$$K(q) \left| \sum_{k=\lceil na \rceil}^{\lfloor nb \rfloor} \left(f\left(\frac{k}{n}\right) - f(x) \right) G_{q,\lambda}(nx - k) \right|. \quad (35)$$

We will use in our estimations the right hand side of the last quantity. For that we need, for $f \in C([a, b], \mathbb{C})$ the first modulus of continuity

$$\omega_1(f, \delta) := \sup_{\substack{x, y \in [a, b] \\ |x - y| \leq \delta}} |f(x) - f(y)|, \quad \delta > 0. \quad (36)$$

The fact $f \in C([a, b], \mathbb{C})$ is equivalent to $\lim_{\delta \rightarrow 0} \omega_1(f, \delta) = 0$, see [8].

3. MAIN RESULTS

We present \mathbb{C} -valued neural network high order approximations to a function given with rates. We start with a trigonometric approximation.

Theorem 3.1. *Let $f \in C^3([a, b], \mathbb{C})$, $0 < \alpha < 1$, $n \in \mathbb{N} : n^{1-\alpha} > 2$, $x \in [a, b]$; denote $\delta := \frac{(b-a)^2}{2} \min\{1, \frac{b-a}{3}\}$. Then*

1)

$$\begin{aligned} |L_n(f, x) - f(x)| \leq K(q) & \left[|f'(x)| \left(\frac{1}{n^\alpha} + (b-a) \gamma A^{-\lambda(n^{1-\alpha}-2)} \right) \right. \\ & + \frac{|f''(x)|}{2} \left(\frac{1}{n^{2\alpha}} + (b-a)^2 \gamma A^{-\lambda(n^{1-\alpha}-2)} \right) + \\ & \left. \left(\frac{\omega_1(f'' + f, \frac{1}{n^\alpha})}{2n^{2\alpha}} + \|f''' + f'\|_\infty \delta \gamma A^{-\lambda(n^{1-\alpha}-2)} \right) \right], \end{aligned} \quad (37)$$

2) if $f'(x) = f''(x) = 0$, we obtain

$$|L_n(f, x) - f(x)| \leq K(q) \left[\frac{\omega_1(f'' + f, \frac{1}{n^\alpha})}{2n^{2\alpha}} + \|f''' + f'\|_\infty \delta \gamma A^{-\lambda(n^{1-\alpha}-2)} \right], \quad (38)$$

notice here the high rate of convergence at $n^{-3\alpha}$,

3) furthermore we get

$$\begin{aligned} \|L_n f - f\|_\infty \leq K(q) & \left[\|f'\|_\infty \left(\frac{1}{n^\alpha} + (b-a) \gamma A^{-\lambda(n^{1-\alpha}-2)} \right) \right. \\ & + \frac{\|f''\|_\infty}{2} \left(\frac{1}{n^{2\alpha}} + (b-a)^2 \gamma A^{-\lambda(n^{1-\alpha}-2)} \right) + \\ & \left. \left(\frac{\omega_1(f'' + f, \frac{1}{n^\alpha})}{2n^{2\alpha}} + \|f''' + f'\|_\infty \delta \gamma A^{-\lambda(n^{1-\alpha}-2)} \right) \right], \end{aligned} \quad (39)$$

i.e. $\lim_{n \rightarrow +\infty} L_n(f) = f$, pointwise and uniformly,

4) and finally, it holds

$$\begin{aligned} \left| L_n(f, x) - f'(x) L_n(\sin(\cdot - x), x) - 2f''(x) L_n\left(\sin^2\left(\frac{\cdot - x}{2}\right), x\right) - f(x) \right| \leq \\ K(q) \left[\frac{\omega_1(f'' + f, \frac{1}{n^\alpha})}{2n^{2\alpha}} + \|f''' + f'\|_\infty \delta \gamma A^{-\lambda(n^{1-\alpha}-2)} \right], \end{aligned} \quad (40)$$

again here we achieve high speed of convergence at $n^{-3\alpha}$.

Proof. Here $f \in C^3([a, b], \mathbb{C})$, and we apply the trigonometric Taylor's formula for $f \in C^2([a, b], \mathbb{C})$, see Theorem 6 of [11].

Let $\frac{k}{n}, x \in [a, b]$, then

$$\begin{aligned} f\left(\frac{k}{n}\right) &= f(x) + f'(x) \sin\left(\frac{k}{n} - x\right) + 2f''(x) \sin^2\left(\frac{\frac{k}{n} - x}{2}\right) + \\ & \int_x^{\frac{k}{n}} [(f''(t) + f(t)) - (f''(x) + f(x))] \sin\left(\frac{k}{n} - t\right) dt. \end{aligned} \quad (41)$$

Hence it holds

$$\begin{aligned} f\left(\frac{k}{n}\right) G_{q,\lambda}(nx-k) &= f(x) G_{q,\lambda}(nx-k) + \\ f'(x) \sin\left(\frac{k}{n}-x\right) G_{q,\lambda}(nx-k) &+ 2f''(x) \sin^2\left(\frac{\frac{k}{n}-x}{2}\right) G_{q,\lambda}(nx-k) + \\ G_{q,\lambda}(nx-k) \left(\int_x^{\frac{k}{n}} [(f''(t)+f(t)) - (f''(x)+f(x))] \sin\left(\frac{k}{n}-t\right) dt\right). \end{aligned} \quad (42)$$

So that we have

$$\begin{aligned} \sum_{k=\lceil na \rceil}^{\lfloor nb \rfloor} f\left(\frac{k}{n}\right) G_{q,\lambda}(nx-k) - f(x) \sum_{k=\lceil na \rceil}^{\lfloor nb \rfloor} G_{q,\lambda}(nx-k) = \\ f'(x) \sum_{k=\lceil na \rceil}^{\lfloor nb \rfloor} G_{q,\lambda}(nx-k) \sin\left(\frac{k}{n}-x\right) + 2f''(x) \sum_{k=\lceil na \rceil}^{\lfloor nb \rfloor} G_{q,\lambda}(nx-k) \sin^2\left(\frac{\frac{k}{n}-x}{2}\right) + \\ \sum_{k=\lceil na \rceil}^{\lfloor nb \rfloor} G_{q,\lambda}(nx-k) \left(\int_x^{\frac{k}{n}} [(f''(t)+f(t)) - (f''(x)+f(x))] \sin\left(\frac{k}{n}-t\right) dt\right). \end{aligned} \quad (43)$$

Thus, we obtain

$$\begin{aligned} L_n^*(f, x) - f(x) \sum_{k=\lceil na \rceil}^{\lfloor nb \rfloor} G_{q,\lambda}(nx-k) = \\ f'(x) L_n^*(\sin(\cdot-x), x) + 2f''(x) L_n^*\left(\sin^2\left(\frac{\cdot-x}{2}\right), x\right) + \Lambda_n(x), \end{aligned} \quad (44)$$

where

$$\Lambda_n(x) := \sum_{k=\lceil na \rceil}^{\lfloor nb \rfloor} G_{q,\lambda}(nx-k) \left(\int_x^{\frac{k}{n}} [(f''(t)+f(t)) - (f''(x)+f(x))] \sin\left(\frac{k}{n}-t\right) dt\right). \quad (45)$$

We call

$$R_2(n) := \int_x^{\frac{k}{n}} [(f''(t)+f(t)) - (f''(x)+f(x))] \sin\left(\frac{k}{n}-t\right) dt. \quad (46)$$

We assume that $b-a > \frac{1}{n^\alpha}$, which is always the case for large enough $n \in \mathbb{N}$, that is when $n > \left[(b-a)^{-\frac{1}{\alpha}}\right]$.

Thus $\left|\frac{k}{n}-x\right| \leq \frac{1}{n^\alpha}$ or $\left|\frac{k}{n}-x\right| > \frac{1}{n^\alpha}$.

In case of $\left|\frac{k}{n}-x\right| \leq \frac{1}{n^\alpha}$, we have the following cases:

i) if $\frac{k}{n} \geq x$, then

$$\begin{aligned} |R_2(n)| &= \left| \int_x^{\frac{k}{n}} [(f''(t)+f(t)) - (f''(x)+f(x))] \sin\left(\frac{k}{n}-t\right) dt \right| \leq \\ &\int_x^{\frac{k}{n}} \omega_1(f''+f, t-x) \left| \sin\left(\frac{k}{n}-t\right) \right| dt \leq \end{aligned} \quad (47)$$

(by $|\sin x| \leq |x|$, $\forall x \in \mathbb{R}$)

$$\begin{aligned} \int_x^{\frac{k}{n}} \omega_1(f'' + f, t-x) \left(\frac{k}{n} - t\right) dt &\leq \omega_1\left(f'' + f, \frac{k}{n} - x\right) \frac{\left(\frac{k}{n} - x\right)^2}{2} \\ &\leq \frac{\omega_1\left(f'' + f, \frac{1}{n^\alpha}\right)}{2n^{2\alpha}}, \end{aligned}$$

that is

$$|R_2(n)| \leq \frac{\omega_1\left(f'' + f, \frac{1}{n^\alpha}\right)}{2n^{2\alpha}}. \quad (48)$$

ii) if $\frac{k}{n} < x$, then

$$\begin{aligned} |R_2(n)| &= \left| \int_x^{\frac{k}{n}} [(f''(t) + f(t)) - (f''(x) + f(x))] \sin\left(\frac{k}{n} - t\right) dt \right| = \\ &\quad \left| \int_{\frac{k}{n}}^x [(f''(t) + f(t)) - (f''(x) + f(x))] \sin\left(\frac{k}{n} - t\right) dt \right| \leq \\ &\quad \int_{\frac{k}{n}}^x |(f''(t) + f(t)) - (f''(x) + f(x))| \left| \sin\left(\frac{k}{n} - t\right) \right| dt \leq \\ &\quad \int_{\frac{k}{n}}^x \omega_1\left(f'' + f, x - \frac{k}{n}\right) \left(t - \frac{k}{n}\right) dt \leq \omega_1\left(f'' + f, x - \frac{k}{n}\right) \frac{\left(x - \frac{k}{n}\right)^2}{2} \\ &\quad \leq \frac{\omega_1\left(f'' + f, \frac{1}{n^\alpha}\right)}{2n^{2\alpha}}. \end{aligned} \quad (49)$$

That is

$$|R_2(n)| \leq \frac{\omega_1\left(f'' + f, \frac{1}{n^\alpha}\right)}{2n^{2\alpha}}. \quad (50)$$

So, we have proved when $|\frac{k}{n} - x| \leq \frac{1}{n^\alpha}$, always it holds

$$|R_2(n)| \leq \frac{\omega_1\left(f'' + f, \frac{1}{n^\alpha}\right)}{2n^{2\alpha}}. \quad (51)$$

Next assume again $\frac{k}{n} \geq x$, then

$$\begin{aligned} |R_2(n)| &= \left| \int_x^{\frac{k}{n}} [(f''(t) + f(t)) - (f''(x) + f(x))] \sin\left(\frac{k}{n} - t\right) dt \right| \leq \\ &\quad \int_x^{\frac{k}{n}} |(f''(t) + f(t)) - (f''(x) + f(x))| \left| \sin\left(\frac{k}{n} - t\right) \right| dt \leq \end{aligned}$$

(by [14], p. 3, abstract mean value theorem, and $|\sin x| \leq 1$ along with $|\sin x| \leq |x|$, $\forall x \in \mathbb{R}$)

$$\begin{aligned} \|f''' + f'\|_\infty \min \left\{ \int_x^{\frac{k}{n}} (t-x) dt, \int_x^{\frac{k}{n}} \left(\frac{k}{n} - t\right) (t-x) dt \right\} &= \\ \|f''' + f'\|_\infty \min \left\{ \frac{\left(\frac{k}{n} - x\right)^2}{2}, \frac{\left(\frac{k}{n} - x\right)^3}{6} \right\} &\leq \\ \|f''' + f'\|_\infty \min \left\{ \frac{(b-a)^2}{2}, \frac{(b-a)^3}{6} \right\} &= \end{aligned} \quad (52)$$

$$\|f''' + f'\|_\infty \frac{(b-a)^2}{2} \min \left\{ 1, \frac{(b-a)}{3} \right\}.$$

Hence

$$|R_2(n)| \leq \|f''' + f'\|_\infty \frac{(b-a)^2}{2} \min \left\{ 1, \frac{(b-a)}{3} \right\}. \quad (53)$$

When $\frac{k}{n} < x$, we have

$$\begin{aligned} |R_2(n)| &= \left| \int_x^{\frac{k}{n}} [(f''(t) + f(t)) - (f''(x) + f(x))] \sin \left(\frac{k}{n} - t \right) dt \right| = \\ &\left| \int_{\frac{k}{n}}^x [(f''(x) + f(x)) - (f''(t) + f(t))] \sin \left(\frac{k}{n} - t \right) dt \right| \leq \\ &\int_{\frac{k}{n}}^x |(f''(x) + f(x)) - (f''(t) + f(t))| \left| \sin \left(\frac{k}{n} - t \right) \right| dt \leq \end{aligned}$$

(as in (52))

$$\begin{aligned} &\|f''' + f'\|_\infty \min \left\{ \int_{\frac{k}{n}}^x (x-t) dt, \int_{\frac{k}{n}}^x (x-t) \left(t - \frac{k}{n} \right) dt \right\} = \\ &\|f''' + f'\|_\infty \min \left\{ \frac{(x - \frac{k}{n})^2}{2}, \frac{(x - \frac{k}{n})^3}{6} \right\} \leq \\ &\|f''' + f'\|_\infty \min \left\{ \frac{(b-a)^2}{2}, \frac{(b-a)^3}{6} \right\} = \\ &\|f''' + f'\|_\infty \frac{(b-a)^2}{2} \min \left\{ 1, \frac{(b-a)}{3} \right\}. \end{aligned} \quad (54)$$

Therefore, always it holds

$$|R_2(n)| \leq \|f''' + f'\|_\infty \frac{(b-a)^2}{2} \min \left\{ 1, \frac{(b-a)}{3} \right\}. \quad (55)$$

And we have

$$\begin{aligned} \Lambda_n(x) &= \sum_{k=\lceil na \rceil}^{\lfloor nb \rfloor} G_{q,\lambda}(nx-k) R_2(n) + \\ &\left\{ \begin{array}{l} k = \lceil na \rceil \\ : \left| \frac{k}{n} - x \right| \leq \frac{1}{n^\alpha} \end{array} \right. \\ &\quad \sum_{k=\lceil na \rceil}^{\lfloor nb \rfloor} G_{q,\lambda}(nx-k) R_2(n). \\ &\left\{ \begin{array}{l} k = \lceil na \rceil \\ : \left| \frac{k}{n} - x \right| > \frac{1}{n^\alpha} \end{array} \right. \end{aligned} \quad (56)$$

Hence it holds

$$\begin{aligned} |\Lambda_n(x)| &\leq \sum_{k=\lceil na \rceil}^{\lfloor nb \rfloor} G_{q,\lambda}(nx-k) |R_2(n)| + \\ &\left\{ \begin{array}{l} k = \lceil na \rceil \\ : \left| \frac{k}{n} - x \right| \leq \frac{1}{n^\alpha} \end{array} \right. \end{aligned}$$

$$\begin{aligned}
& \sum_{k=\lceil na \rceil}^{\lfloor nb \rfloor} G_{q,\lambda}(nx-k) |R_2(n)| \leq \tag{57} \\
& \left\{ \begin{array}{l} k = \lceil na \rceil \\ : \left| \frac{k}{n} - x \right| > \frac{1}{n^\alpha} \end{array} \right. \\
& \left(\sum_{k=\lceil na \rceil}^{\lfloor nb \rfloor} G_{q,\lambda}(nx-k) \right) \frac{\omega_1\left(f'' + f, \frac{1}{n^\alpha}\right)}{2n^{2\alpha}} + \\
& \sum_{k=\lceil na \rceil}^{\lfloor nb \rfloor} G_{q,\lambda}(nx-k) \|f''' + f'\|_\infty \frac{(b-a)^2}{2} \min\left\{1, \frac{(b-a)}{3}\right\} \stackrel{\text{(by (30))}}{\leq} \\
& \left\{ \begin{array}{l} k = \lceil na \rceil \\ : \left| \frac{k}{n} - x \right| > \frac{1}{n^\alpha} \end{array} \right. \\
& \frac{\omega_1\left(f'' + f, \frac{1}{n^\alpha}\right)}{2n^{2\alpha}} + \|f''' + f'\|_\infty \frac{(b-a)^2}{2} \min\left\{1, \frac{(b-a)}{3}\right\} \\
& \left(\sum_{k=\lceil na \rceil}^{\lfloor nb \rfloor} G_{q,\lambda}(nx-k) \right) \stackrel{\text{(by Theorem 2.3)}}{\leq} \tag{58} \\
& \frac{\omega_1\left(f'' + f, \frac{1}{n^\alpha}\right)}{2n^{2\alpha}} + \|f''' + f'\|_\infty \frac{(b-a)^2}{2} \min\left\{1, \frac{(b-a)}{3}\right\} \gamma A^{-\lambda(n^{1-\alpha}-2)}.
\end{aligned}$$

From now on for convenience we call

$$\delta := \frac{(b-a)^2}{2} \min\left\{1, \frac{(b-a)}{3}\right\}. \tag{59}$$

Consequently, we have derived that

$$|\Lambda_n(x)| \leq \frac{\omega_1\left(f'' + f, \frac{1}{n^\alpha}\right)}{2n^{2\alpha}} + \|f''' + f'\|_\infty \delta \gamma A^{-\lambda(n^{1-\alpha}-2)}. \tag{60}$$

Next we use again $|\sin x| \leq |x|$, $\forall x \in \mathbb{R}$.

We have that

$$L_n^*(\sin(\cdot - x), x) = \sum_{k=\lceil na \rceil}^{\lfloor nb \rfloor} G_{q,\lambda}(nx-k) \sin\left(\frac{k}{n} - x\right), \tag{61}$$

and

$$\begin{aligned}
|L_n^*(\sin(\cdot - x), x)| & \leq \sum_{k=\lceil na \rceil}^{\lfloor nb \rfloor} G_{q,\lambda}(nx-k) \left| \sin\left(\frac{k}{n} - x\right) \right| = \\
& \sum_{k=\lceil na \rceil}^{\lfloor nb \rfloor} G_{q,\lambda}(nx-k) \left| \sin\left(\frac{k}{n} - x\right) \right| + \\
& \left\{ \begin{array}{l} k = \lceil na \rceil \\ : \left| \frac{k}{n} - x \right| \leq \frac{1}{n^\alpha} \end{array} \right.
\end{aligned}$$

$$\sum_{\substack{k = [na] \\ : \left| \frac{k}{n} - x \right| > \frac{1}{n^\alpha}}}^{[nb]} G_{q,\lambda}(nx - k) \left| \sin \left(\frac{k}{n} - x \right) \right| \leq \quad (62)$$

$$\sum_{\substack{k = [na] \\ : \left| \frac{k}{n} - x \right| \leq \frac{1}{n^\alpha}}}^{[nb]} G_{q,\lambda}(nx - k) \left| \frac{k}{n} - x \right| +$$

$$\sum_{\substack{k = [na] \\ : \left| \frac{k}{n} - x \right| > \frac{1}{n^\alpha}}}^{[nb]} G_{q,\lambda}(nx - k) \left| \frac{k}{n} - x \right| \leq$$

$$\frac{1}{n^\alpha} + (b - a) \left(\sum_{\substack{k = [na] \\ : \left| \frac{k}{n} - x \right| > \frac{1}{n^\alpha}}}^{[nb]} G_{q,\lambda}(nx - k) \right) \stackrel{\text{(by (27))}}{\leq} \quad (63)$$

$$\frac{1}{n^\alpha} + (b - a) \gamma A^{-\lambda(n^{1-\alpha}-2)}.$$

We found that

$$|L_n^*(\sin(\cdot - x), x)| \leq \frac{1}{n^\alpha} + (b - a) \gamma A^{-\lambda(n^{1-\alpha}-2)}. \quad (64)$$

Next we estimate

$$L_n^* \left(\sin^2 \left(\frac{\cdot - x}{2} \right), x \right) = \sum_{k=[na]}^{[nb]} G_{q,\lambda}(nx - k) \sin^2 \left(\frac{\frac{k}{n} - x}{2} \right), \quad (65)$$

We have that (by $|\sin x| \leq |x|, \forall x \in \mathbb{R}$)

$$L_n^* \left(\sin^2 \left(\frac{\cdot - x}{2} \right), x \right) = \sum_{k=[na]}^{[nb]} G_{q,\lambda}(nx - k) \left| \sin \left(\frac{\frac{k}{n} - x}{2} \right) \right|^2 \leq$$

$$\frac{1}{4} \sum_{k=[na]}^{[nb]} G_{q,\lambda}(nx - k) \left| \frac{k}{n} - x \right|^2 =$$

$$\frac{1}{4} \left[\sum_{\substack{k = [na] \\ : \left| \frac{k}{n} - x \right| \leq \frac{1}{n^\alpha}}}^{[nb]} G_{q,\lambda}(nx - k) \left| \frac{k}{n} - x \right|^2 + \right.$$

$$\left\{ \begin{array}{l} \sum_{k=\lceil na \rceil}^{\lfloor nb \rfloor} G_{q,\lambda}(nx-k) \left| \frac{k}{n} - x \right|^2 \\ : \left| \frac{k}{n} - x \right| > \frac{1}{n^\alpha} \end{array} \right\} \leq \quad (66)$$

$$\frac{1}{4} \left[\frac{1}{n^{2\alpha}} + (b-a)^2 \gamma A^{-\lambda(n^{1-\alpha}-2)} \right].$$

That is

$$L_n^* \left(\sin^2 \left(\frac{\cdot - x}{2} \right), x \right) \leq \frac{1}{4} \left[\frac{1}{n^{2\alpha}} + (b-a)^2 \gamma A^{-\lambda(n^{1-\alpha}-2)} \right]. \quad (67)$$

Consequently we have derived:

1)

$$\begin{aligned} |L_n(f, x) - f(x)| &\leq K(q) \left[|f'(x)| \left(\frac{1}{n^\alpha} + (b-a) \gamma A^{-\lambda(n^{1-\alpha}-2)} \right) \right. \\ &\quad \left. + \frac{|f''(x)|}{2} \left(\frac{1}{n^{2\alpha}} + (b-a)^2 \gamma A^{-\lambda(n^{1-\alpha}-2)} \right) + \right. \\ &\quad \left. \left(\frac{\omega_1(f'' + f, \frac{1}{n^\alpha})}{2n^{2\alpha}} + \|f''' + f'\|_\infty \delta \gamma A^{-\lambda(n^{1-\alpha}-2)} \right) \right]. \end{aligned} \quad (68)$$

2) if $f'(x) = f''(x) = 0$, by (68), we obtain

$$|L_n(f, x) - f(x)| \leq K(q) \left[\frac{\omega_1(f'' + f, \frac{1}{n^\alpha})}{2n^{2\alpha}} + \|f''' + f'\|_\infty \delta \gamma A^{-\lambda(n^{1-\alpha}-2)} \right], \quad (69)$$

notice here the high rate of convergence at $n^{-3\alpha}$.

3) Furthermore, by (68), we get

$$\begin{aligned} \|L_n f - f\|_\infty &\leq K(q) \left[\|f'\|_\infty \left(\frac{1}{n^\alpha} + (b-a) \gamma A^{-\lambda(n^{1-\alpha}-2)} \right) + \right. \\ &\quad \left. \frac{\|f''\|_\infty}{2} \left(\frac{1}{n^{2\alpha}} + (b-a)^2 \gamma A^{-\lambda(n^{1-\alpha}-2)} \right) + \right. \\ &\quad \left. \left(\frac{\omega_1(f'' + f, \frac{1}{n^\alpha})}{2n^{2\alpha}} + \|f''' + f'\|_\infty \delta \gamma A^{-\lambda(n^{1-\alpha}-2)} \right) \right]. \end{aligned} \quad (70)$$

We derive that $\lim_{n \rightarrow +\infty} L_n(f) = f$, pointwise and uniformly.

We observe that

$$\begin{aligned} L_n(f, x) - f'(x) L_n(\sin(\cdot - x), x) - 2f''(x) L_n \left(\sin^2 \left(\frac{\cdot - x}{2} \right), x \right) - f(x) = \\ \frac{L_n^*(f, x)}{\sum_{k=\lceil na \rceil}^{\lfloor nb \rfloor} G_{q,\lambda}(nx-k)} - f'(x) \frac{L_n^*(\sin(\cdot - x), x)}{\sum_{k=\lceil na \rceil}^{\lfloor nb \rfloor} G_{q,\lambda}(nx-k)} - \\ 2f''(x) \frac{L_n^*(\sin^2(\frac{\cdot - x}{2}), x)}{\sum_{k=\lceil na \rceil}^{\lfloor nb \rfloor} G_{q,\lambda}(nx-k)} - f(x) \left(\frac{\sum_{k=\lceil na \rceil}^{\lfloor nb \rfloor} G_{q,\lambda}(nx-k)}{\sum_{k=\lceil na \rceil}^{\lfloor nb \rfloor} G_{q,\lambda}(nx-k)} \right) = \end{aligned} \quad (71)$$

$$\frac{1}{\sum_{k=\lceil na \rceil}^{\lfloor nb \rfloor} G_{q,\lambda}(nx-k)} [L_n^*(f, x) - f'(x) L_n^*(\sin(\cdot - x), x) - 2f''(x) L_n^*\left(\sin^2\left(\frac{\cdot - x}{2}\right), x\right) - f(x) \sum_{k=\lceil na \rceil}^{\lfloor nb \rfloor} G_{q,\lambda}(nx-k)] = \frac{1}{\sum_{k=\lceil na \rceil}^{\lfloor nb \rfloor} G_{q,\lambda}(nx-k)} (\Lambda_n(x)). \quad (72)$$

Finally, we obtain ($\forall x \in [a, b]$, $n \in \mathbb{N}$):

$$\begin{aligned} & 4) \left| L_n(f, x) - f'(x) L_n(\sin(\cdot - x), x) - 2f''(x) L_n\left(\sin^2\left(\frac{\cdot - x}{2}\right), x\right) - f(x) \right| \stackrel{(28)}{\leq} \\ & K(q) |\Lambda_n(x)| \stackrel{(60)}{\leq} \\ & K(q) \left[\frac{\omega_1(f'' + f, \frac{1}{n^\alpha})}{2n^{2\alpha}} + \|f''' + f'\|_\infty \delta \gamma A^{-\lambda(n^{1-\alpha}-2)} \right]. \quad (73) \end{aligned}$$

The theorem is proved. \square

We continue with a hyperbolic high order neural network approximation.

Theorem 3.2. *Let $f \in C^3([a, b], \mathbb{C})$, $0 < \alpha < 1$, $n \in \mathbb{N} : n^{1-\alpha} > 2$, $x \in [a, b]$. Then*

1)

$$\begin{aligned} |L_n(f, x) - f(x)| & \leq K(q) \cosh(b-a) \left[|f'(x)| \left(\frac{1}{n^\alpha} + (b-a) \gamma A^{-\lambda(n^{1-\alpha}-2)} \right) \right. \\ & \quad \left. \frac{|f''(x)|}{2} \left(\frac{1}{n^{2\alpha}} + (b-a)^2 \gamma A^{-\lambda(n^{1-\alpha}-2)} \right) + \right. \\ & \quad \left. \frac{1}{2} \left(\frac{\omega_1(f'' - f, \frac{1}{n^\alpha})}{n^{2\alpha}} + \frac{\|f''' - f'\|_\infty (b-a)^3}{3} \gamma A^{-\lambda(n^{1-\alpha}-2)} \right) \right], \quad (74) \end{aligned}$$

2) if $f'(x) = f''(x) = 0$, we obtain

$$\begin{aligned} |L_n(f, x) - f(x)| & \leq \frac{K(q) \cosh(b-a)}{2} \\ & \left[\frac{\omega_1(f'' - f, \frac{1}{n^\alpha})}{n^{2\alpha}} + \frac{\|f''' - f'\|_\infty (b-a)^3}{3} \gamma A^{-\lambda(n^{1-\alpha}-2)} \right], \quad (75) \end{aligned}$$

notice here the high rate of convergence at $n^{-3\alpha}$,

3) furthermore, we get

$$\begin{aligned} \|L_n f - f\|_\infty & \leq K(q) \cosh(b-a) \left[\|f'\|_\infty \left(\frac{1}{n^\alpha} + (b-a) \gamma A^{-\lambda(n^{1-\alpha}-2)} \right) \right. \\ & \quad \left. \frac{\|f''\|_\infty}{2} \left(\frac{1}{n^{2\alpha}} + (b-a)^2 \gamma A^{-\lambda(n^{1-\alpha}-2)} \right) + \right. \\ & \quad \left. \frac{1}{2} \left(\frac{\omega_1(f'' - f, \frac{1}{n^\alpha})}{n^{2\alpha}} + \frac{\|f''' - f'\|_\infty (b-a)^3}{3} \gamma A^{-\lambda(n^{1-\alpha}-2)} \right) \right], \quad (76) \end{aligned}$$

it follows that $\lim_{n \rightarrow +\infty} L_n(f) = f$, pointwise and uniformly,

and
4)

$$\left| L_n(f, x) - f'(x) L_n(\sinh(\cdot - x), x) - 2f''(x) L_n\left(\sinh^2\left(\frac{\cdot - x}{2}\right), x\right) - f(x) \right| \leq \frac{K(q) \cosh(b-a)}{2} \left[\frac{\omega_1(f'' - f, \frac{1}{n^\alpha})}{n^{2\alpha}} + \frac{\|f''' - f'\|_\infty (b-a)^3}{3} \gamma A^{-\lambda(n^{1-\alpha}-2)} \right], \quad (77)$$

again here we achieve high speed of convergence at $n^{-3\alpha}$.

Proof. By the mean value theorem we have that

$$\sinh x = \sinh x - \sinh 0 = (\cosh \xi)(x - 0),$$

for some ξ between $\{0, x\}$, for any $x \in \mathbb{R}$.

Hence

$$|\sinh x| \leq \|\cosh\|_{\infty, [-(b-a), b-a]} |x|, \quad \forall x \in [-(b-a), b-a]. \quad (78)$$

That is, there exists $M \geq 1$ such that

$$|\sinh x| \leq M|x|, \quad \forall x \in [-(b-a), b-a], \quad (79)$$

where $M := \|\cosh\|_{\infty, [-(b-a), b-a]} = \cosh(b-a)$.

Here $f \in C^3([a, b], \mathbb{C})$, and we apply the hyperbolic Taylor's formula for $f \in C^2([a, b], \mathbb{C})$, see Theorem 7 of [11].

Let $\frac{k}{n}, x \in [a, b]$, then

$$\begin{aligned} f\left(\frac{k}{n}\right) &= f(x) + f'(x) \sinh\left(\frac{k}{n} - x\right) + 2f''(x) \sinh^2\left(\frac{\frac{k}{n} - x}{2}\right) + \\ &\int_x^{\frac{k}{n}} [(f''(t) - f(t)) - (f''(x) - f(x))] \sinh\left(\frac{k}{n} - t\right) dt. \end{aligned} \quad (80)$$

Hence it holds

$$\begin{aligned} f\left(\frac{k}{n}\right) G_{q,\lambda}(nx - k) &= f(x) G_{q,\lambda}(nx - k) + \\ f'(x) \sinh\left(\frac{k}{n} - x\right) G_{q,\lambda}(nx - k) &+ 2f''(x) \sinh^2\left(\frac{\frac{k}{n} - x}{2}\right) G_{q,\lambda}(nx - k) + \\ G_{q,\lambda}(nx - k) \left(\int_x^{\frac{k}{n}} [(f''(t) - f(t)) - (f''(x) - f(x))] \sinh\left(\frac{k}{n} - t\right) dt \right). \end{aligned} \quad (81)$$

So that we have

$$\begin{aligned} \sum_{k=\lceil na \rceil}^{\lfloor nb \rfloor} f\left(\frac{k}{n}\right) G_{q,\lambda}(nx - k) - f(x) \sum_{k=\lceil na \rceil}^{\lfloor nb \rfloor} G_{q,\lambda}(nx - k) &= \\ f'(x) \sum_{k=\lceil na \rceil}^{\lfloor nb \rfloor} G_{q,\lambda}(nx - k) \sinh\left(\frac{k}{n} - x\right) + 2f''(x) \sum_{k=\lceil na \rceil}^{\lfloor nb \rfloor} G_{q,\lambda}(nx - k) \sinh^2\left(\frac{\frac{k}{n} - x}{2}\right) &+ \\ \sum_{k=\lceil na \rceil}^{\lfloor nb \rfloor} G_{q,\lambda}(nx - k) \left(\int_x^{\frac{k}{n}} [(f''(t) - f(t)) - (f''(x) - f(x))] \sinh\left(\frac{k}{n} - t\right) dt \right). \end{aligned} \quad (82)$$

Thus, we obtain

$$L_n^*(f, x) - f(x) \sum_{k=\lceil na \rceil}^{\lfloor nb \rfloor} G_{q,\lambda}(nx - k) = f'(x) L_n^*(\sinh(\cdot - x), x) + 2f''(x) L_n^*\left(\sinh^2\left(\frac{\cdot - x}{2}\right), x\right) + \Lambda_n(x), \quad (83)$$

where

$$\Lambda_n(x) := \sum_{k=\lceil na \rceil}^{\lfloor nb \rfloor} G_{q,\lambda}(nx - k) \left(\int_x^{\frac{k}{n}} [(f''(t) - f(t)) - (f''(x) - f(x))] \sinh\left(\frac{k}{n} - t\right) dt \right). \quad (84)$$

We call

$$R_2(n) := \int_x^{\frac{k}{n}} [(f''(t) - f(t)) - (f''(x) - f(x))] \sinh\left(\frac{k}{n} - t\right) dt. \quad (85)$$

We assume that $b - a > \frac{1}{n^\alpha}$, which is always the case for large enough $n \in \mathbb{N}$, that is when $n > \left[(b - a)^{-\frac{1}{\alpha}}\right]$.

Thus $\left|\frac{k}{n} - x\right| \leq \frac{1}{n^\alpha}$ or $\left|\frac{k}{n} - x\right| > \frac{1}{n^\alpha}$.

In case of $\left|\frac{k}{n} - x\right| \leq \frac{1}{n^\alpha}$, we have the following cases:

i) if $\frac{k}{n} \geq x$, then

$$\begin{aligned} |R_2(n)| &= \left| \int_x^{\frac{k}{n}} [(f''(t) - f(t)) - (f''(x) - f(x))] \sinh\left(\frac{k}{n} - t\right) dt \right| \leq \\ &\int_x^{\frac{k}{n}} \omega_1(f'' - f, t - x) \left| \sinh\left(\frac{k}{n} - t\right) \right| dt \stackrel{(79)}{\leq} \\ &\int_x^{\frac{k}{n}} \omega_1(f'' - f, t - x) M\left(\frac{k}{n} - t\right) dt \leq M\omega_1\left(f'' - f, \frac{k}{n} - x\right) \int_x^{\frac{k}{n}} \left(\frac{k}{n} - t\right) dt = \\ &M\omega_1\left(f'' - f, \frac{k}{n} - x\right) \frac{\left(\frac{k}{n} - x\right)^2}{2} \leq \frac{M\omega_1\left(f'' - f, \frac{1}{n^\alpha}\right)}{2n^{2\alpha}}, \end{aligned} \quad (86)$$

that is

$$|R_2(n)| \leq \frac{M\omega_1\left(f'' - f, \frac{1}{n^\alpha}\right)}{2n^{2\alpha}}. \quad (87)$$

ii) if $\frac{k}{n} < x$, then

$$\begin{aligned} |R_2(n)| &= \left| \int_x^{\frac{k}{n}} [(f''(t) - f(t)) - (f''(x) - f(x))] \sinh\left(\frac{k}{n} - t\right) dt \right| = \\ &\left| \int_{\frac{k}{n}}^x [(f''(t) - f(t)) - (f''(x) - f(x))] \sinh\left(\frac{k}{n} - t\right) dt \right| \leq \\ &\int_{\frac{k}{n}}^x |(f''(t) - f(t)) - (f''(x) - f(x))| \left| \sinh\left(\frac{k}{n} - t\right) \right| dt \leq \\ &M\omega_1\left(f'' - f, x - \frac{k}{n}\right) \int_{\frac{k}{n}}^x \left(t - \frac{k}{n}\right) dt = M\omega_1\left(f'' - f, x - \frac{k}{n}\right) \frac{\left(x - \frac{k}{n}\right)^2}{2} \end{aligned} \quad (88)$$

$$\leq \frac{M\omega_1\left(f'' - f, \frac{1}{n^\alpha}\right)}{2n^{2\alpha}},$$

that is

$$|R_2(n)| \leq \frac{M\omega_1\left(f'' - f, \frac{1}{n^\alpha}\right)}{2n^{2\alpha}}. \quad (89)$$

So, we have proved when $|\frac{k}{n} - x| \leq \frac{1}{n^\alpha}$, always it holds

$$|R_2(n)| \leq \frac{M\omega_1\left(f'' - f, \frac{1}{n^\alpha}\right)}{2n^{2\alpha}}. \quad (90)$$

Next assume again $\frac{k}{n} \geq x$, then

$$\begin{aligned} |R_2(n)| &= \left| \int_x^{\frac{k}{n}} [(f''(t) - f(t)) - (f''(x) - f(x))] \sinh\left(\frac{k}{n} - t\right) dt \right| \leq \\ &\int_x^{\frac{k}{n}} |(f''(t) - f(t)) - (f''(x) - f(x))| \left| \sinh\left(\frac{k}{n} - t\right) \right| dt \leq \end{aligned}$$

(by [14], p. 3, abstract mean value theorem)

$$\begin{aligned} M \|f''' - f'\|_\infty \int_x^{\frac{k}{n}} \left(\frac{k}{n} - t\right) (t - x) dt &= \quad (91) \\ M \|f''' - f'\|_\infty \frac{\left(\frac{k}{n} - x\right)^3}{6} &\leq M \|f''' - f'\|_\infty \frac{(b-a)^3}{6}. \end{aligned}$$

Hence

$$|R_2(n)| \leq \frac{M \|f''' - f'\|_\infty (b-a)^3}{6}. \quad (92)$$

When $\frac{k}{n} < x$, we have

$$\begin{aligned} |R_2(n)| &= \left| \int_{\frac{k}{n}}^x [(f''(t) - f(t)) - (f''(x) - f(x))] \sinh\left(\frac{k}{n} - t\right) dt \right| \leq \\ &\int_{\frac{k}{n}}^x |(f''(t) - f(t)) - (f''(x) - f(x))| \left| \sinh\left(\frac{k}{n} - t\right) \right| dt \leq \end{aligned}$$

(as in (91))

$$\begin{aligned} M \|f''' - f'\|_\infty \int_{\frac{k}{n}}^x (x - t) \left(t - \frac{k}{n}\right) dt &= \quad (93) \\ M \|f''' - f'\|_\infty \frac{\left(x - \frac{k}{n}\right)^3}{6} &\leq M \|f''' - f'\|_\infty \frac{(b-a)^3}{6}. \end{aligned}$$

Therefore, always it holds

$$|R_2(n)| \leq \frac{M \|f''' - f'\|_\infty (b-a)^3}{6}. \quad (94)$$

And we have

$$\begin{aligned} \Lambda_n(x) &= \sum_{k=0}^{\lfloor nb \rfloor} G_{q,\lambda}(nx - k) R_2(n) + \\ &\begin{cases} k = \lceil na \rceil \\ : \left| \frac{k}{n} - x \right| \leq \frac{1}{n^\alpha} \end{cases} \quad (95) \end{aligned}$$

$$\sum_{k=\lceil na \rceil}^{\lfloor nb \rfloor} G_{q,\lambda}(nx-k) R_2(n).$$

$$\left\{ \begin{array}{l} k = \lceil na \rceil \\ : \left| \frac{k}{n} - x \right| > \frac{1}{n^\alpha} \end{array} \right.$$

Hence it holds

$$|\Lambda_n(x)| \leq \sum_{k=\lceil na \rceil}^{\lfloor nb \rfloor} G_{q,\lambda}(nx-k) |R_2(n)| + \tag{96}$$

$$\left\{ \begin{array}{l} k = \lceil na \rceil \\ : \left| \frac{k}{n} - x \right| \leq \frac{1}{n^\alpha} \end{array} \right.$$

$$\sum_{k=\lceil na \rceil}^{\lfloor nb \rfloor} G_{q,\lambda}(nx-k) |R_2(n)| \leq$$

$$\left\{ \begin{array}{l} k = \lceil na \rceil \\ : \left| \frac{k}{n} - x \right| > \frac{1}{n^\alpha} \end{array} \right.$$

$$\left(\sum_{k=\lceil na \rceil}^{\lfloor nb \rfloor} G_{q,\lambda}(nx-k) \right) \frac{\omega_1(f'' - f, \frac{1}{n^\alpha}) M}{2n^{2\alpha}} +$$

$$\left(\sum_{k=\lceil na \rceil}^{\lfloor nb \rfloor} G_{q,\lambda}(nx-k) \right) \frac{M \|f''' - f'\|_\infty (b-a)^3}{6} \stackrel{((30))}{\leq}$$

$$\frac{M \omega_1(f'' - f, \frac{1}{n^\alpha})}{2n^{2\alpha}} + \frac{M \|f''' - f'\|_\infty (b-a)^3}{6}$$

$$\left(\sum_{k=\lceil na \rceil}^{\lfloor nb \rfloor} G_{q,\lambda}(nx-k) \right) \stackrel{\text{(by Theorem 2.3)}}{\leq} \tag{97}$$

$$M \frac{\omega_1(f'' - f, \frac{1}{n^\alpha})}{2n^{2\alpha}} + \frac{M \|f''' - f'\|_\infty (b-a)^3}{6} \gamma A^{-\lambda(n^{1-\alpha}-2)}.$$

Consequently, we have derived that

$$|\Lambda_n(x)| \leq \frac{M}{2} \left[\frac{\omega_1(f'' - f, \frac{1}{n^\alpha})}{n^{2\alpha}} + \frac{\|f''' - f'\|_\infty (b-a)^3}{3} \gamma A^{-\lambda(n^{1-\alpha}-2)} \right]. \tag{98}$$

We have that

$$L_n^*(\sinh(\cdot - x), x) = \sum_{k=\lceil na \rceil}^{\lfloor nb \rfloor} G_{q,\lambda}(nx-k) \sinh\left(\frac{k}{n} - x\right), \tag{99}$$

and

$$\begin{aligned}
|L_n^*(\sinh(\cdot - x), x)| &\leq \sum_{k=\lceil na \rceil}^{\lfloor nb \rfloor} G_{q,\lambda}(nx - k) \left| \sinh\left(\frac{k}{n} - x\right) \right| = \\
&\quad \sum_{\substack{k=\lceil na \rceil \\ : \left| \frac{k}{n} - x \right| \leq \frac{1}{n^\alpha}}}^{\lfloor nb \rfloor} G_{q,\lambda}(nx - k) \left| \sinh\left(\frac{k}{n} - x\right) \right| + \\
&\quad \sum_{\substack{k=\lceil na \rceil \\ : \left| \frac{k}{n} - x \right| > \frac{1}{n^\alpha}}}^{\lfloor nb \rfloor} G_{q,\lambda}(nx - k) \left| \sinh\left(\frac{k}{n} - x\right) \right| \leq \quad (100)
\end{aligned}$$

$$\begin{aligned}
&M \left[\sum_{\substack{k=\lceil na \rceil \\ : \left| \frac{k}{n} - x \right| \leq \frac{1}{n^\alpha}}}^{\lfloor nb \rfloor} G_{q,\lambda}(nx - k) \left| \frac{k}{n} - x \right| + \right. \\
&\quad \left. \sum_{\substack{k=\lceil na \rceil \\ : \left| \frac{k}{n} - x \right| > \frac{1}{n^\alpha}}}^{\lfloor nb \rfloor} G_{q,\lambda}(nx - k) \left| \frac{k}{n} - x \right| \right] \leq \quad (101) \\
&M \left[\frac{1}{n^\alpha} + (b - a) \left(\sum_{\substack{k=\lceil na \rceil \\ : \left| \frac{k}{n} - x \right| > \frac{1}{n^\alpha}}}^{\lfloor nb \rfloor} G_{q,\lambda}(nx - k) \right) \right] \stackrel{\text{(by (27))}}{\leq} \\
&M \left[\frac{1}{n^\alpha} + (b - a) \gamma A^{-\lambda(n^{1-\alpha}-2)} \right].
\end{aligned}$$

We found that

$$|L_n^*(\sinh(\cdot - x), x)| \leq M \left[\frac{1}{n^\alpha} + (b - a) \gamma A^{-\lambda(n^{1-\alpha}-2)} \right]. \quad (102)$$

Next we estimate

$$L_n^* \left(\sinh^2 \left(\frac{\cdot - x}{2} \right), x \right) = \sum_{k=\lceil na \rceil}^{\lfloor nb \rfloor} G_{q,\lambda}(nx - k) \sinh^2 \left(\frac{k}{n} - x \right), \quad (103)$$

We have that

$$L_n^* \left(\sinh^2 \left(\frac{\cdot - x}{2} \right), x \right) = \sum_{k=\lceil na \rceil}^{\lfloor nb \rfloor} G_{q,\lambda}(nx - k) \left| \sinh \left(\frac{k}{n} - x \right) \right|^2 \leq$$

$$\begin{aligned} & \frac{M}{4} \sum_{k=\lceil na \rceil}^{\lfloor nb \rfloor} G_{q,\lambda}(nx-k) \left| \frac{k}{n} - x \right|^2 = \tag{104} \\ & \frac{M}{4} \left[\sum_{\substack{k=\lceil na \rceil \\ : \left| \frac{k}{n} - x \right| \leq \frac{1}{n^\alpha}}}^{\lfloor nb \rfloor} G_{q,\lambda}(nx-k) \left| \frac{k}{n} - x \right|^2 + \right. \\ & \left. \sum_{\substack{k=\lceil na \rceil \\ : \left| \frac{k}{n} - x \right| > \frac{1}{n^\alpha}}}^{\lfloor nb \rfloor} G_{q,\lambda}(nx-k) \left| \frac{k}{n} - x \right|^2 \right] \leq \\ & \frac{M}{4} \left[\frac{1}{n^{2\alpha}} + (b-a)^2 \gamma A^{-\lambda(n^{1-\alpha}-2)} \right]. \end{aligned}$$

That is

$$L_n^* \left(\sinh^2 \left(\frac{\cdot - x}{2} \right), x \right) \leq \frac{M}{4} \left[\frac{1}{n^{2\alpha}} + (b-a)^2 \gamma A^{-\lambda(n^{1-\alpha}-2)} \right]. \tag{105}$$

By (35) and putting together (83), (98), (102) and (105) we derive 1)

$$\begin{aligned} |L_n(f, x) - f(x)| & \leq K(q) M \left[|f'(x)| \left(\frac{1}{n^\alpha} + (b-a) \gamma A^{-\lambda(n^{1-\alpha}-2)} \right) + \right. \\ & \quad \left. \frac{|f''(x)|}{2} \left(\frac{1}{n^{2\alpha}} + (b-a)^2 \gamma A^{-\lambda(n^{1-\alpha}-2)} \right) + \right. \\ & \quad \left. \frac{1}{2} \left(\frac{\omega_1(f'' - f, \frac{1}{n^\alpha})}{n^{2\alpha}} + \frac{\|f''' - f'\|_\infty (b-a)^3}{3} \gamma A^{-\lambda(n^{1-\alpha}-2)} \right) \right]. \tag{106} \end{aligned}$$

2) If $f'(x) = f''(x) = 0$, by (106), we obtain

$$\begin{aligned} |L_n(f, x) - f(x)| & \leq \\ & \frac{MK(q)}{2} \left[\frac{\omega_1(f'' - f, \frac{1}{n^\alpha})}{n^{2\alpha}} + \frac{\|f''' - f'\|_\infty (b-a)^3}{3} \gamma A^{-\lambda(n^{1-\alpha}-2)} \right], \tag{107} \end{aligned}$$

notice here the high rate of convergence at $n^{-3\alpha}$.

3) Furthermore, by (106), we get

$$\begin{aligned} \|L_n f - f\|_\infty & \leq K(q) M \left[\|f'\|_\infty \left(\frac{1}{n^\alpha} + (b-a) \gamma A^{-\lambda(n^{1-\alpha}-2)} \right) \right. \\ & \quad \left. \frac{\|f''\|_\infty}{2} \left(\frac{1}{n^{2\alpha}} + (b-a)^2 \gamma A^{-\lambda(n^{1-\alpha}-2)} \right) + \right. \\ & \quad \left. \frac{1}{2} \left(\frac{\omega_1(f'' - f, \frac{1}{n^\alpha})}{n^{2\alpha}} + \frac{\|f''' - f'\|_\infty (b-a)^3}{3} \gamma A^{-\lambda(n^{1-\alpha}-2)} \right) \right]. \tag{108} \end{aligned}$$

It follows that $\lim_{n \rightarrow +\infty} L_n(f) = f$, pointwise and uniformly.

We observe that

$$L_n(f, x) - f'(x) L_n(\sinh(\cdot - x), x) - 2f''(x) L_n\left(\sinh^2\left(\frac{\cdot - x}{2}\right), x\right) - f(x) =$$

$$\frac{L_n^*(f, x)}{\sum_{k=\lceil na \rceil}^{\lfloor nb \rfloor} G_{q,\lambda}(nx - k)} - f'(x) \frac{L_n^*(\sinh(\cdot - x), x)}{\sum_{k=\lceil na \rceil}^{\lfloor nb \rfloor} G_{q,\lambda}(nx - k)} -$$

$$2f''(x) \frac{L_n^*\left(\sinh^2\left(\frac{\cdot - x}{2}\right), x\right)}{\sum_{k=\lceil na \rceil}^{\lfloor nb \rfloor} G_{q,\lambda}(nx - k)} - f(x) \left(\frac{\sum_{k=\lceil na \rceil}^{\lfloor nb \rfloor} G_{q,\lambda}(nx - k)}{\sum_{k=\lceil na \rceil}^{\lfloor nb \rfloor} G_{q,\lambda}(nx - k)} \right) = \quad (109)$$

$$\frac{1}{\sum_{k=\lceil na \rceil}^{\lfloor nb \rfloor} G_{q,\lambda}(nx - k)} [L_n^*(f, x) - f'(x) L_n^*(\sinh(\cdot - x), x) -$$

$$2f''(x) L_n^*\left(\sinh^2\left(\frac{\cdot - x}{2}\right), x\right) - f(x) \sum_{k=\lceil na \rceil}^{\lfloor nb \rfloor} G_{q,\lambda}(nx - k)] = \quad (110)$$

$$\frac{1}{\sum_{k=\lceil na \rceil}^{\lfloor nb \rfloor} G_{q,\lambda}(nx - k)} (\Lambda_n(x)).$$

Finally, we obtain ($\forall x \in [a, b]$, $n \in \mathbb{N}$):

4)

$$\left| L_n(f, x) - f'(x) L_n(\sinh(\cdot - x), x) - 2f''(x) L_n\left(\sinh^2\left(\frac{\cdot - x}{2}\right), x\right) - f(x) \right| \stackrel{(28)}{\leq}$$

$$K(q) |\Lambda_n(x)| \stackrel{(98)}{\leq}$$

$$\frac{K(q) M}{2} \left[\frac{\omega_1\left(f'' - f, \frac{1}{n^\alpha}\right)}{n^{2\alpha}} + \frac{\|f''' - f'\|_\infty (b-a)^3}{3} \gamma A^{-\lambda(n^{1-\alpha}-2)} \right]. \quad (111)$$

The theorem is established. \square

Next follows a mixed hyperbolic-trigonometric high order neural network approximation.

Theorem 3.3. *Let $f \in C^5([a, b], \mathbb{C})$, $0 < \alpha < 1$, $n \in \mathbb{N} : n^{1-\alpha} > 2$, $x \in [a, b]$. Then*

1)

$$\left| L_n(f, x) - f(x) - \frac{f'(x)}{2} L_n((\sinh(\cdot - x) + \sin(\cdot - x)), x) \right.$$

$$\left. - \frac{f''(x)}{2} L_n((\cosh(\cdot - x) - \cos(\cdot - x)), x) \right. \quad (112)$$

$$\left. - \frac{f^{(3)}(x)}{2} L_n((\sinh(\cdot - x) - \sin(\cdot - x)), x) \right.$$

$$\left. - f^{(4)}(x) L_n\left(\left(\sinh^2\left(\frac{\cdot - x}{2}\right) - \sin^2\left(\frac{\cdot - x}{2}\right)\right), x\right) \right| \leq$$

$$\frac{K(q) (\cosh(b-a) + 1)}{4}$$

$$\left[\frac{\omega_1\left(f^{(4)} - f, \frac{1}{n^\alpha}\right)}{n^{2\alpha}} + \frac{\|f^{(5)} - f'\|_\infty (b-a)^3}{3} \gamma A^{-\lambda(n^{1-\alpha}-2)} \right], \quad (113)$$

2) if $f^{(i)}(x) = 0$, $i = 1, 2, 3, 4$, we get

$$|L_n(f, x) - f(x)| \leq \frac{K(q)(\cosh(b-a) + 1)}{4} \left[\frac{\omega_1(f^{(4)} - f, \frac{1}{n^\alpha})}{n^{2\alpha}} + \frac{\|f^{(5)} - f'\|_\infty (b-a)^3}{3} \gamma A^{-\lambda(n^{1-\alpha}-2)} \right], \quad (114)$$

in the last (114) observe the high speed of convergence at $n^{-3\alpha}$.

Proof. Here $f \in C^5([a, b], \mathbb{C})$, and we apply the hyperbolic-trigonometric Taylor's formula for $f \in C^4([a, b], \mathbb{C})$, see Theorem 8 of [11].

Let $\frac{k}{n}, x \in [a, b]$, then

$$\begin{aligned} & f\left(\frac{k}{n}\right) - f(x) - f'(x) \left(\frac{\sinh\left(\frac{k}{n} - x\right) + \sin\left(\frac{k}{n} - x\right)}{2} \right) \\ & - f''(x) \left(\frac{\cosh\left(\frac{k}{n} - x\right) - \cos\left(\frac{k}{n} - x\right)}{2} \right) \\ & - f^{(3)}(x) \left(\frac{\sinh\left(\frac{k}{n} - x\right) - \sin\left(\frac{k}{n} - x\right)}{2} \right) \\ & - f^{(4)}(x) \left(\sinh^2\left(\frac{\frac{k}{n} - x}{2}\right) - \sin^2\left(\frac{\frac{k}{n} - x}{2}\right) \right) = \quad (115) \\ & \int_x^{\frac{k}{n}} \left[\left(f^{(4)}(t) - f^{(4)}(x) \right) - \left(f^{(4)}(x) - f^{(4)}(x) \right) \right] \left(\frac{\sinh\left(\frac{k}{n} - t\right) - \sin\left(\frac{k}{n} - t\right)}{2} \right) dt \\ & =: R_4\left(\frac{k}{n}, x\right). \end{aligned}$$

As in Theorems 3.1, 3.2 we derive

$$\begin{aligned} & L_n^*(f, x) - f(x) \sum_{k=\lceil na \rceil}^{\lfloor nb \rfloor} G_{q,\lambda}(nx - k) - \\ & \frac{f'(x)}{2} L_n^* \left((\sinh(\cdot - x) + \sin(\cdot - x)), x \right) - \\ & \frac{f''(x)}{2} L_n^* \left((\cosh(\cdot - x) - \cos(\cdot - x)), x \right) - \\ & \frac{f^{(3)}(x)}{2} L_n^* \left((\sinh(\cdot - x) - \sin(\cdot - x)), x \right) - \\ & \frac{f^{(4)}(x)}{2} L_n^* \left(\left(\sinh^2\left(\frac{\cdot - x}{2}\right) - \sin^2\left(\frac{\cdot - x}{2}\right) \right), x \right) = \Phi_n(x), \quad (116) \end{aligned}$$

where

$$\Phi_n(x) := \sum_{k=\lceil na \rceil}^{\lfloor nb \rfloor} G_{q,\lambda}(nx - k) R_4\left(\frac{k}{n}, x\right). \quad (117)$$

Without loss of generality we can assume that $n > \left\lceil (b-a)^{-\frac{1}{\alpha}} \right\rceil$.

Thus $\left| \frac{k}{n} - x \right| \leq \frac{1}{n^\alpha}$ or $\left| \frac{k}{n} - x \right| > \frac{1}{n^\alpha}$.

In case of $\left| \frac{k}{n} - x \right| \leq \frac{1}{n^\alpha}$, we have the following cases:

i) if $\frac{k}{n} \geq x$, then

$$\begin{aligned} & \left| R_4 \left(\frac{k}{n}, x \right) \right| = \\ & \left| \frac{1}{2} \int_x^{\frac{k}{n}} \left[\left(f^{(4)}(t) - f(t) \right) - \left(f^{(4)}(x) - f(x) \right) \right] \left(\sinh \left(\frac{k}{n} - t \right) - \sin \left(\frac{k}{n} - t \right) \right) dt \right| \leq \\ & \frac{1}{2} \int_x^{\frac{k}{n}} \left| \left(f^{(4)}(t) - f(t) \right) - \left(f^{(4)}(x) - f(x) \right) \right| \left| \sinh \left(\frac{k}{n} - t \right) - \sin \left(\frac{k}{n} - t \right) \right| dt \leq \end{aligned} \quad (118)$$

$$\begin{aligned} & \frac{1}{2} \int_x^{\frac{k}{n}} \omega_1 \left(f^{(4)} - f, t - x \right) \left(\left| \sinh \left(\frac{k}{n} - t \right) \right| + \left| \sin \left(\frac{k}{n} - t \right) \right| \right) dt \leq \\ & \frac{\omega_1 \left(f^{(4)} - f, \frac{k}{n} - x \right)}{2} \int_x^{\frac{k}{n}} \left(\cosh(b-a) \left(\frac{k}{n} - t \right) + \left(\frac{k}{n} - t \right) \right) dt = \\ & \frac{(\cosh(b-a) + 1) \omega_1 \left(f^{(4)} - f, \frac{k}{n} - x \right)}{2} \int_x^{\frac{k}{n}} \left(\frac{k}{n} - t \right) dt = \\ & \frac{(\cosh(b-a) + 1) \omega_1 \left(f^{(4)} - f, \frac{k}{n} - x \right)}{4} \left(\frac{k}{n} - x \right)^2 \leq \\ & \frac{(\cosh(b-a) + 1) \omega_1 \left(f^{(4)} - f, \frac{1}{n^\alpha} \right)}{4n^{2\alpha}}. \end{aligned} \quad (119)$$

That is, when $\frac{k}{n} \geq x$, then

$$\left| R_4 \left(\frac{k}{n}, x \right) \right| \leq \frac{(\cosh(b-a) + 1) \omega_1 \left(f^{(4)} - f, \frac{1}{n^\alpha} \right)}{4n^{2\alpha}}. \quad (120)$$

ii) if $\frac{k}{n} < x$, then

$$\begin{aligned} & \left| R_4 \left(\frac{k}{n}, x \right) \right| = \\ & \left| \frac{1}{2} \int_{\frac{k}{n}}^x \left[\left(f^{(4)}(t) - f(t) \right) - \left(f^{(4)}(x) - f(x) \right) \right] \left(\sinh \left(\frac{k}{n} - t \right) - \sin \left(\frac{k}{n} - t \right) \right) dt \right| \leq \\ & \frac{\omega_1 \left(f^{(4)} - f, x - \frac{k}{n} \right)}{2} \int_{\frac{k}{n}}^x \left(\cosh(b-a) \left(t - \frac{k}{n} \right) + \left(t - \frac{k}{n} \right) \right) dt = \\ & \frac{(\cosh(b-a) + 1) \omega_1 \left(f^{(4)} - f, x - \frac{k}{n} \right)}{2} \int_{\frac{k}{n}}^x \left(t - \frac{k}{n} \right) dt = \\ & \frac{(\cosh(b-a) + 1) \omega_1 \left(f^{(4)} - f, x - \frac{k}{n} \right)}{4} \left(x - \frac{k}{n} \right)^2 \leq \\ & \leq \frac{(\cosh(b-a) + 1) \omega_1 \left(f^{(4)} - f, \frac{1}{n^\alpha} \right)}{4n^{2\alpha}}. \end{aligned} \quad (121)$$

Consequently, when $\left| \frac{k}{n} - x \right| \leq \frac{1}{n^\alpha}$, we always obtain that

$$\left| R_4 \left(\frac{k}{n}, x \right) \right| \leq \frac{(\cosh(b-a) + 1) \omega_1 \left(f^{(4)} - f, \frac{1}{n^\alpha} \right)}{4n^{2\alpha}}. \quad (122)$$

Next assume again $\frac{k}{n} \geq x$, then

$$\left| R_4 \left(\frac{k}{n}, x \right) \right| \leq$$

$$\frac{1}{2} \int_x^{\frac{k}{n}} \left| \left(f^{(4)}(t) - f(t) \right) - \left(f^{(4)}(x) - f(x) \right) \right| \left| \sinh \left(\frac{k}{n} - t \right) - \sin \left(\frac{k}{n} - t \right) \right| dt \leq$$

(by [14], p. 3, abstract mean value theorem)

$$\begin{aligned} & \frac{\|f^{(5)} - f'\|_\infty}{2} \int_x^{\frac{k}{n}} (t-x) \left[\cosh(b-a) \left(\frac{k}{n} - t \right) + \left(\frac{k}{n} - t \right) \right] dt = \\ & \frac{\|f^{(5)} - f'\|_\infty}{2} (\cosh(b-a) + 1) \int_x^{\frac{k}{n}} \left(\frac{k}{n} - t \right) (t-x) dt = \\ & \frac{\|f^{(5)} - f'\|_\infty (\cosh(b-a) + 1)}{12} \left(\frac{k}{n} - x \right)^3 \leq \\ & \leq \frac{\|f^{(5)} - f'\|_\infty (\cosh(b-a) + 1) (b-a)^3}{12}. \end{aligned} \tag{123}$$

Hence

$$\left| R_4 \left(\frac{k}{n}, x \right) \right| \leq \frac{\|f^{(5)} - f'\|_\infty (\cosh(b-a) + 1) (b-a)^3}{12}. \tag{124}$$

When $\frac{k}{n} < x$, we have

$$\begin{aligned} & \left| R_4 \left(\frac{k}{n}, x \right) \right| \leq \\ & \frac{1}{2} \int_{\frac{k}{n}}^x \left| \left(f^{(4)}(t) - f(t) \right) - \left(f^{(4)}(x) - f(x) \right) \right| \left| \sinh \left(\frac{k}{n} - t \right) - \sin \left(\frac{k}{n} - t \right) \right| dt \leq \\ & \frac{\|f^{(5)} - f'\|_\infty}{2} \int_{\frac{k}{n}}^x (x-t) \left[\cosh(b-a) \left(t - \frac{k}{n} \right) + \left(t - \frac{k}{n} \right) \right] dt = \\ & \frac{\|f^{(5)} - f'\|_\infty (\cosh(b-a) + 1)}{2} \int_{\frac{k}{n}}^x (x-t) \left(t - \frac{k}{n} \right) dt = \\ & \frac{\|f^{(5)} - f'\|_\infty (\cosh(b-a) + 1)}{12} \left(x - \frac{k}{n} \right)^3 \leq \\ & \leq \frac{\|f^{(5)} - f'\|_\infty (\cosh(b-a) + 1) (b-a)^3}{12}. \end{aligned} \tag{125}$$

So, it is always true that

$$\left| R_4 \left(\frac{k}{n}, x \right) \right| \leq \frac{\|f^{(5)} - f'\|_\infty (\cosh(b-a) + 1) (b-a)^3}{12}. \tag{126}$$

Thus

$$\begin{aligned} |\Phi_n(x)| & \leq \sum_{k=\lceil na \rceil}^{\lfloor nb \rfloor} G_{q,\lambda}(nx-k) \left| R_4 \left(\frac{k}{n}, x \right) \right| = \\ & \sum_{k=\lceil na \rceil}^{\lfloor nb \rfloor} G_{q,\lambda}(nx-k) \left| R_4 \left(\frac{k}{n}, x \right) \right| + \\ & \left\{ \begin{array}{l} k = \lceil na \rceil \\ : \left| \frac{k}{n} - x \right| \leq \frac{1}{n^\alpha} \end{array} \right. \end{aligned} \tag{127}$$

$$\left(\sum_{\substack{k = \lceil na \rceil \\ \left| \frac{k}{n} - x \right| > \frac{1}{n^\alpha}}}^{\lfloor nb \rfloor} G_{q,\lambda}(nx - k) \left| R_4 \left(\frac{k}{n}, x \right) \right| \leq \right. \\ \left. \left(\sum_{\substack{k = \lceil na \rceil \\ \left| \frac{k}{n} - x \right| \leq \frac{1}{n^\alpha}}}^{\lfloor nb \rfloor} G_{q,\lambda}(nx - k) \right) \frac{(\cosh(b-a) + 1) \omega_1 \left(f^{(4)} - f, \frac{1}{n^\alpha} \right)}{4n^{2\alpha}} + \right) \quad (128)$$

$$\left(\sum_{\substack{k = \lceil na \rceil \\ \left| \frac{k}{n} - x \right| > \frac{1}{n^\alpha}}}^{\lfloor nb \rfloor} G_{q,\lambda}(nx - k) \right) \\ \left(\frac{\|f^{(5)} - f'\|_\infty (\cosh(b-a) + 1) (b-a)^3}{12} \right) \stackrel{\text{(by Theorem 2.3)}}{\leq} \\ \frac{(\cosh(b-a) + 1) \omega_1 \left(f^{(4)} - f, \frac{1}{n^\alpha} \right)}{4n^{2\alpha}} + \quad (129) \\ \frac{\|f^{(5)} - f'\|_\infty (\cosh(b-a) + 1) (b-a)^3}{12} \gamma A^{-\lambda(n^{1-\alpha}-2)}.$$

We have proved that

$$|\Phi_n(x)| \leq \frac{(\cosh(b-a) + 1)}{4} \\ \left[\frac{\omega_1 \left(f^{(4)} - f, \frac{1}{n^\alpha} \right)}{n^{2\alpha}} + \frac{\|f^{(5)} - f'\|_\infty (b-a)^3}{3} \gamma A^{-\lambda(n^{1-\alpha}-2)} \right]. \quad (130)$$

We observe that

$$L_n(f, x) - f(x) - \frac{f'(x)}{2} L_n((\sinh(\cdot - x) + \sin(\cdot - x)), x) \\ - \frac{f''(x)}{2} L_n((\cosh(\cdot - x) - \cos(\cdot - x)), x) \\ - \frac{f^{(3)}(x)}{2} L_n((\sinh(\cdot - x) - \sin(\cdot - x)), x) \\ - f^{(4)}(x) L_n\left(\left(\sinh^2\left(\frac{\cdot - x}{2}\right) - \sin^2\left(\frac{\cdot - x}{2}\right)\right), x\right) = \quad (131)$$

$$\left[L_n^*(f, x) - f(x) \sum_{k=\lceil na \rceil}^{\lfloor nb \rfloor} G_{q,\lambda}(nx - k) - \frac{f'(x)}{2} L_n^*((\sinh(\cdot - x) + \sin(\cdot - x)), x) \right. \\ \left. - \frac{f''(x)}{2} L_n^*((\cosh(\cdot - x) - \cos(\cdot - x)), x) \right. \\ \left. - \frac{f^{(3)}(x)}{2} L_n^*((\sinh(\cdot - x) - \sin(\cdot - x)), x) \right. \quad (132)$$

$$-f^{(4)}(x) L_n^* \left(\left(\sinh^2 \left(\frac{\cdot - x}{2} \right) - \sin^2 \left(\frac{\cdot - x}{2} \right) \right), x \right) \frac{1}{\sum_{k=\lceil na \rceil}^{\lfloor nb \rfloor} G_{q,\lambda}(nx - k)}$$

$$= \frac{\Phi_n(x)}{\sum_{k=\lceil na \rceil}^{\lfloor nb \rfloor} G_{q,\lambda}(nx - k)}.$$

Finally, we obtain ($\forall x \in [a, b], n \in \mathbb{N}$):

$$\left| L_n(f, x) - f(x) - \frac{f'(x)}{2} L_n((\sinh(\cdot - x) + \sin(\cdot - x)), x) \right.$$

$$\left. - \frac{f''(x)}{2} L_n((\cosh(\cdot - x) - \cos(\cdot - x)), x) \right.$$

$$\left. - \frac{f^{(3)}(x)}{2} L_n((\sinh(\cdot - x) - \sin(\cdot - x)), x) \right.$$

$$\left. - f^{(4)}(x) L_n \left(\left(\sinh^2 \left(\frac{\cdot - x}{2} \right) - \sin^2 \left(\frac{\cdot - x}{2} \right) \right), x \right) \right| =$$

$$\frac{|\Phi_n(x)|}{\sum_{k=\lceil na \rceil}^{\lfloor nb \rfloor} G_{q,\lambda}(nx - k)} \leq K(q) |\Phi_n(x)| \stackrel{(by(130))}{\leq} \frac{K(q) (\cosh(b - a) + 1)}{4}$$

$$\left[\frac{\omega_1(f^{(4)} - f, \frac{1}{n^\alpha})}{n^{2\alpha}} + \frac{\|f^{(5)} - f'\|_\infty (b - a)^3}{3} \gamma A^{-\lambda(n^{1-\alpha} - 2)} \right]. \quad (134)$$

The theorem is proved. \square

We continue with a general trigonometric result.

Theorem 3.4. *Let $f \in C^5([a, b], \mathbb{C})$, $0 < \alpha < 1$, $n \in \mathbb{N} : n^{1-\alpha} > 2$, $x \in [a, b]$. Let also $\bar{\alpha}, \bar{\beta} \in \mathbb{R}$ with $\bar{\alpha}\bar{\beta}(\bar{\alpha}^2 - \bar{\beta}^2) \neq 0$. Then*

1)

$$\left| L_n(f, x) - f(x) - \frac{f'(x)}{\bar{\alpha}\bar{\beta}(\bar{\beta}^2 - \bar{\alpha}^2)} L_n \left(\left(\bar{\beta}^3 \sin(\bar{\alpha}(\cdot - x)) - \bar{\alpha}^3 \sin(\bar{\beta}(\cdot - x)) \right), x \right) \right.$$

$$\left. - \frac{f''(x)}{(\bar{\beta}^2 - \bar{\alpha}^2)} L_n((\cos(\bar{\alpha}(\cdot - x)) - \cos(\bar{\beta}(\cdot - x))), x) \right.$$

$$\left. - \frac{f'''(x)}{\bar{\alpha}\bar{\beta}(\bar{\beta}^2 - \bar{\alpha}^2)} L_n((\bar{\beta} \sin(\bar{\alpha}(\cdot - x)) - \bar{\alpha} \sin(\bar{\beta}(\cdot - x))), x) \right.$$

$$\left. - \left(\frac{2f^{(4)}(x) + (\bar{\alpha}^2 + \bar{\beta}^2)f''(x)}{(\bar{\alpha}\bar{\beta})^2(\bar{\beta}^2 - \bar{\alpha}^2)} \right) \right.$$

$$\left. L_n \left(\left(\bar{\beta}^2 \sin^2 \left(\frac{\bar{\alpha}(\cdot - x)}{2} \right) - \bar{\alpha}^2 \sin^2 \left(\frac{\bar{\beta}(\cdot - x)}{2} \right) \right), x \right) \right| \leq \quad (135)$$

$$\frac{K(q)}{|\bar{\beta}^2 - \bar{\alpha}^2|} \left[\frac{\omega_1 \left(\left(f^{(4)} + (\bar{\alpha}^2 + \bar{\beta}^2)f'' + \bar{\alpha}^2\bar{\beta}^2 f \right), \frac{1}{n^\alpha} \right)}{n^{2\alpha}} + \right.$$

$$\left[\frac{\|f^{(5)} + (\bar{\alpha}^2 + \bar{\beta}^2) f''' + \bar{\alpha}^2 \bar{\beta}^2 f'\|_{\infty} (b-a)^3}{3} \gamma A^{-\lambda(n^{1-\alpha}-2)} \right],$$

2) if $f^{(i)}(x) = 0$, $i = 1, 2, 3, 4$, we get

$$\begin{aligned} |L_n(f, x) - f(x)| &\leq \frac{K(q)}{|\bar{\beta}^2 - \bar{\alpha}^2|} \\ &\left[\frac{\omega_1\left(\left(f^{(4)} + (\bar{\alpha}^2 + \bar{\beta}^2) f'' + \bar{\alpha}^2 \bar{\beta}^2 f\right), \frac{1}{n^\alpha}\right)}{n^{2\alpha}} + \right. \\ &\left. \frac{\|f^{(5)} + (\bar{\alpha}^2 + \bar{\beta}^2) f''' + \bar{\alpha}^2 \bar{\beta}^2 f'\|_{\infty} (b-a)^3}{3} \gamma A^{-\lambda(n^{1-\alpha}-2)} \right]. \end{aligned} \quad (136)$$

The high speed of convergence in (1) and (2) is $n^{-3\alpha}$.

Proof. As similar to Theorem 3.3 is omitted. It is based on Theorem 9 of [11]. \square

We finish with a general hyperbolic result.

Theorem 3.5. Let $f \in C^5([a, b], \mathbb{C})$, $0 < \alpha < 1$, $n \in \mathbb{N} : n^{1-\alpha} > 2$, $x \in [a, b]$. Let also $\bar{\alpha}, \bar{\beta} \in \mathbb{R}$ with $\bar{\alpha}\bar{\beta}(\bar{\alpha}^2 - \bar{\beta}^2) \neq 0$. Then

1)

$$\begin{aligned} \left| L_n(f, x) - f(x) - \frac{f'(x)}{\bar{\alpha}\bar{\beta}(\bar{\beta}^2 - \bar{\alpha}^2)} L_n\left(\left(\bar{\beta}^3 \sinh(\bar{\alpha}(\cdot - x)) - \bar{\alpha}^3 \sinh(\bar{\beta}(\cdot - x))\right), x\right) \right. \\ \left. - \frac{f''(x)}{\bar{\beta}^2 - \bar{\alpha}^2} L_n\left(\left(\cosh(\bar{\beta}(\cdot - x)) - \cosh(\bar{\alpha}(\cdot - x))\right), x\right) \right. \\ \left. - \frac{f'''(x)}{\bar{\alpha}\bar{\beta}(\bar{\beta}^2 - \bar{\alpha}^2)} L_n\left(\left(\bar{\alpha} \sinh(\bar{\beta}(\cdot - x)) - \bar{\beta} \sinh(\bar{\alpha}(\cdot - x))\right), x\right) \right. \\ \left. - \left(\frac{2(f^{(4)}(x) - (\bar{\alpha}^2 + \bar{\beta}^2) f''(x))}{(\bar{\alpha}\bar{\beta})^2(\bar{\beta}^2 - \bar{\alpha}^2)}\right) \right. \\ \left. L_n\left(\left(\bar{\alpha}^2 \sinh^2\left(\frac{\bar{\beta}(\cdot - x)}{2}\right) - \bar{\beta}^2 \sinh^2\left(\frac{\bar{\alpha}(\cdot - x)}{2}\right)\right), x\right) \right| \leq \quad (137) \\ \frac{K(q) \cosh(b-a)}{|\bar{\beta}^2 - \bar{\alpha}^2|} \left[\frac{\omega_1\left(\left(f^{(4)} - (\bar{\alpha}^2 + \bar{\beta}^2) f'' + \bar{\alpha}^2 \bar{\beta}^2 f\right), \frac{1}{n^\alpha}\right)}{n^{2\alpha}} + \right. \\ \left. \frac{\|f^{(5)} - (\bar{\alpha}^2 + \bar{\beta}^2) f''' + \bar{\alpha}^2 \bar{\beta}^2 f'\|_{\infty} (b-a)^3}{3} \gamma A^{-\lambda(n^{1-\alpha}-2)} \right], \end{aligned}$$

2) if $f^{(i)}(x) = 0$, $i = 1, 2, 3, 4$, we get

$$|L_n(f, x) - f(x)| \leq \frac{K(q) \cosh(b-a)}{|\bar{\beta}^2 - \bar{\alpha}^2|} \quad (138)$$

$$\left[\frac{\omega_1 \left(\left(f^{(4)} - (\bar{\alpha}^2 + \bar{\beta}^2) f'' + \bar{\alpha}^2 \bar{\beta}^2 f \right), \frac{1}{n^\alpha} \right)}{n^{2\alpha}} + \frac{\left\| f^{(5)} - (\bar{\alpha}^2 + \bar{\beta}^2) f''' + \bar{\alpha}^2 \bar{\beta}^2 f' \right\|_\infty (b-a)^3}{3} \gamma_A^{-\lambda(n^{1-\alpha}-2)} \right].$$

The high speed of convergence in (1) and (2) is $n^{-3\alpha}$.

Proof. As similar to Theorem 3.3 is omitted. It is based on Theorem 10 of [11]. \square

REFERENCES

- [1] G.A. Anastassiou, *Rate of convergence of some neural network operators to the unit-univariate case*, J. Math. Anal. Appl. 212 (1997), 237-262.
- [2] G.A. Anastassiou, *Quantitative Approximations*, Chapman & Hall / CRC, Boca Raton, New York, 2001.
- [3] G.A. Anastassiou, *Univariate hyperbolic tangent neural network approximation*, Mathematics and Computer Modelling, 53 (2011), 1111-1132.
- [4] G.A. Anastassiou, *Multivariate hyperbolic tangent neural network approximation*, Computers and Mathematics, 61 (2011), 809-821.
- [5] G.A. Anastassiou, *Multivariate sigmoidal neural network approximation*, Neural Networks, 24 (2011), 378-386.
- [6] G.A. Anastassiou, *Intelligent Systems: Approximation by Artificial Neural Networks*, Intelligent Systems Reference Library, Vol. 19, Springer, Heidelberg, 2011.
- [7] G.A. Anastassiou, *Univariate sigmoidal neural network approximation*, J. of Computational Analysis and Applications, Vol. 14, No. 4, 2012, 659-690.
- [8] G.A. Anastassiou, *Vector fractional Korovkin type Approximations*, Dynamic Systems and Applications, 26(2017), 81-104.
- [9] G.A. Anastassiou, *General sigmoid based Banach space valued neural network approximation*, J. of Computational Analysis and Applications, 31(4) (2023), 520-534.
- [10] G.A. Anastassiou, *q -Deformed and λ -parametrized A -generalized logistic function induced Banach space valued multivariate multi layer neural network approximations*, submitted, 2022.
- [11] G.A. Anastassiou, *Opial and Ostrowski type inequalities based on trigonometric and hyperbolic type Taylor formulae*, submitted, 2023.
- [12] Z. Chen and F. Cao, *The approximation operators with sigmoidal functions*, Computers and Mathematics with Applications, 58 (2009), 758-765.
- [13] S. Haykin, *Neural Networks: A Comprehensive Foundation* (2 ed.), Prentice Hall, New York, 1998.
- [14] V. Lakshmikantham, G.E. Ladas, *Differential Equations in Abstract Spaces*, Academic Press, San Diego, New York, 1972.
- [15] W. McCulloch and W. Pitts, *A logical calculus of the ideas immanent in nervous activity*, Bulletin of Mathematical Biophysics, 7 (1943), 115-133.
- [16] T.M. Mitchell, *Machine Learning*, WCB-McGraw-Hill, New York, 1997.

GEORGE A. ANASTASSIOU
 DEPARTMENT OF MATHEMATICAL SCIENCES, UNIVERSITY OF MEMPHIS,
 MEMPHIS, TN 38152, U.S.A.
E-mail address: ganastss@memphis.edu